


# 2015

## Organic Production and IPM Guide for Snap Beans



NYS IPM Publication No. 132



Cornell University  
Cooperative Extension



Integrated Pest Management



New York State  
Department of  
Agriculture & Markets

# 2015 PRODUCTION GUIDE FOR ORGANIC BEANS FOR PROCESSING

---

## Coordinating Editor

Abby Seaman\* (NYS IPM Program)

## Contributors and Resources

George Abawi (Cornell University, Section of Plant Pathology and Plant-Microbe Biology, retired)

Thomas Bjorkman (Cornell University, Vegetable Crop Physiology)

Ann Cobb (Cornell University, Section of Plant Pathology and Plant-Microbe Biology, retired)

Helene Dillard (Dean of the College of Agricultural and Environmental Sciences at UC Davis, formerly Cornell University, Section of Plant Pathology and Plant-Microbe Biology)

Vern Grubinger (University of Vermont, Vegetable and Berry Specialist)

Beth Gugino (The Pennsylvania State University, Department of Plant Pathology)

Robert Hadad (Cornell Vegetable Program)

Julie Kikkert (Cornell Vegetable Program)

Michael Helms\* (Cornell University, Pesticide Management Education Program)

Margaret T. McGrath (Cornell University, Section of Plant Pathology and Plant-Microbe Biology)

Charles L. Mohler (Cornell University, Section of Soil and Crop Sciences, retired)

Brian Nault\* (Cornell University, Department of Entomology)

Anu Rangarajan (Local Economies Project, The New World foundation, formerly Cornell University Small Farms Program, Section of Horticulture)

Thomas A. Zitter (Cornell University, Section of Plant Pathology and Plant-Microbe Biology, retired)

*\*Pesticide Information and Regulatory Compliance*

## Staff Writers

Mary Kirkwyland and Elizabeth Thomas (New York State IPM Program)

## Editing for the 2015 update

Mary Kirkwyland (Cornell University, NYSAES, New York State IPM Program)

## Special Appreciation

Format based on the Integrated Crop and Pest Management Guidelines for Commercial Vegetable Production (Reference 4). Content Editors Stephen Reiners, Curtis H. Petzoldt, and Abby Seaman with numerous Discipline Editors.

## Funded in part by the New York State Department of Agriculture and Markets

The information in this guide reflects the current authors' best effort to interpret a complex body of scientific research, and to translate this into practical management options. Following the guidance provided in this guide does not assure compliance with any applicable law, rule, regulation or standard, or the achievement of particular discharge levels from agricultural land.

Every effort has been made to provide correct, complete, and up-to-date pest management information for New York State at the time this publication was released for printing (June 2015). Changes in pesticide registrations and regulations, occurring after publication are available in county Cornell Cooperative Extension offices or from the Pesticide Management Education Program web site (<http://pmep.cce.cornell.edu>). Trade names used herein are for convenience only. No endorsement of products is intended, nor is criticism of unnamed products implied.

***This guide is not a substitute for pesticide labeling. Always read the product label before applying any pesticide.***

Updates and additions to this guide are available at [http://www.nysipm.cornell.edu/organic\\_guide](http://www.nysipm.cornell.edu/organic_guide). Please submit comments or suggested changes for these guides to [organicguides@gmail.com](mailto:organicguides@gmail.com).

How to cite this publication: Seaman, Abby, Editor. (2015). Production Guide for Organic Beans for Processing. Publisher: New York State Integrated Pest Management Program, Cornell University (New York State Agricultural Experiment Station, Geneva, NY). 50 pages.

<b>INTRODUCTION .....</b>	<b>1</b>
<b>1. General Organic Management Practices .....</b>	<b>1</b>
<b>2. Soil Health .....</b>	<b>1</b>
<b>3. Cover Crops .....</b>	<b>2</b>
3.1 Goals and Timing for Cover Crops .....	2
3.2 Legumes Cover Crops .....	2
3.3 Non-Legume Cover Crops .....	2
3.4 Biofumigant Cover Crops .....	2
<b>4. Field Selection .....</b>	<b>4</b>
4.1 Certifying Requirements .....	4
4.2 Crop Rotation Plan .....	4
4.3 Pest History .....	5
4.4 Soil and Air Drainage .....	5
<b>5. Weed Management .....</b>	<b>5</b>
5.1 Record Keeping .....	6
5.2 Weed Management Methods .....	6
<b>6. Recommended Varieties .....</b>	<b>6</b>
<b>7. Planting Methods .....</b>	<b>7</b>
<b>8. Crop &amp; Soil Nutrient Management .....</b>	<b>7</b>
8.1 Fertility .....	8
8.2 Preparing an Organic Nutrient Budget .....	8
<b>9. Harvesting .....</b>	<b>11</b>
<b>10. Using Organic Pesticides .....</b>	<b>11</b>
10.1 Sprayer Calibration and Application .....	11
10.2 Regulatory Considerations .....	11
10.3 Optimizing Pesticide Effectiveness .....	12
<b>11. Disease Management .....</b>	<b>12</b>
11.1 Sclerotinia White Mold, <i>Sclerotinia sclerotiorum</i> .....	15
11.2 Botrytis Gray Mold, <i>Botrytis cinerea</i> .....	17
11.3 Pod-flecking complex (PFC), <i>Alternaria alternata</i> and <i>Plectosporium tabacinum</i> .....	19
11.4 Root Rot and Damping-Off .....	20
11.5 Bacterial Diseases .....	23
11.6 Virus Diseases .....	25
11.7 Bean Rust, <i>Uromyces appendiculatus</i> .....	26
<b>12. Root-Lesion Nematode, <i>Pratylenchus penetrans</i> .....</b>	<b>28</b>
<b>13. Insect Management .....</b>	<b>28</b>
13.1 Mexican Bean Beetle .....	31
13.2 Potato Leafhopper (PLH), <i>Empoasca fabae</i> .....	34
13.3 Seedcorn Maggot, <i>Delia platura</i> .....	36
13.4 European Corn Borer (ECB), <i>Ostrinia nubilalis</i> .....	37
13.5 Two-Spotted Spider Mite, <i>Tetranychus urticae</i> .....	39
13.6 Tarnished Plant Bug (TPB), <i>Lygus lineolaris</i> .....	41
<b>14. Slugs .....</b>	<b>43</b>
<b>15. Pesticides &amp; Abbreviations Mentioned in this Publication .....</b>	<b>44</b>
<b>16. References .....</b>	<b>46</b>

## INTRODUCTION

This guide for organic production of snap beans provides an outline of cultural and pest management practices and includes topics that have an impact on improving plant health and reducing pest problems. It is divided into sections, but the interrelated quality of organic cropping systems makes each section relevant to the others.

The guide attempts to compile the most current information available, but acknowledges that effective means of control are not available for some pests. More research on growing crops organically is needed, especially in the area of pest management. Future revisions will incorporate new information, providing organic growers with a complete set of useful practices to help them achieve success.

This guide uses the term Integrated Pest Management (IPM), which like organic production, emphasizes cultural, biological, and mechanical practices to minimize pest outbreaks. With limited pest control products available for use in many organic production systems, an integrated approach to pest management is essential. IPM techniques such as identifying and assessing pest populations, keeping accurate pest history records, selecting the proper site, and preventing pest outbreaks through use of crop rotation, resistant varieties and biological controls are important to producing a high quality crop.

## 1. GENERAL ORGANIC MANAGEMENT PRACTICES

### 1.1 Organic Certification

To use a certified organic label, farming operations that gross more than \$5,000 per year in organic products must be certified by a U.S. Department of Agriculture National Organic Program (NOP) accredited certifying agency. The choice of certifier may be dictated by the processor or by the target market. [A list of accredited certifiers](#) (Link 4) operating in New York can be found on the New York State Department of Agriculture and Markets [Organic Farming Development/Assistance](#) web page (Link 5). See more certification and regulatory details under Section 4.1: *Certification Requirements* and Section 10: *Using Organic Pesticides*.

### 1.2 Organic Farm Plan

An organic farm plan is central to the certification process. The farm plan describes production, handling, and record-keeping systems, and demonstrates to certifiers an understanding of organic practices for a specific crop. The process of developing the plan can be very valuable in terms of anticipating potential issues and challenges, and fosters thinking of the farm as a whole system. Soil, nutrient, pest, and weed management are all interrelated on organic farms and must be managed in concert for success.

Certifying organizations may be able to provide a template for the farm plan. The following description of the farm plan is from the NOP web site:

“The Organic Food Production Act of 1990 (OFPA or Act) requires that all crop, wild crop, livestock, and handling operations requiring certification submit an organic system plan to their certifying agent and, where applicable, the State Organic Program (SOP). The organic system plan is a detailed description of how an operation will achieve, document, and sustain compliance with all applicable provisions in the OFPA and these regulations. The certifying agent must concur that the proposed organic system plan fulfills the requirements of subpart C, and any subsequent modification of the organic plan by the producer or handler must receive the approval of the certifying agent.”

More details may be found at the Agricultural Marketing Service’s [National Organic Program website](#) (Link 6). The [National Sustainable Agriculture Information Service](#), (formerly ATTRA), has produced a guide to organic certification that includes templates for developing an organic farm plan (Link 7). The [Rodale Institute](#) has also developed resources for transitioning to organic and developing an organic farm plan (Link 8).

## 2. SOIL HEALTH

Healthy soil is the basis of organic farming. Regular additions of organic matter in the form of cover crops, compost, or manure create a soil that is biologically active, with good structure and capacity to hold nutrients and water (note that any raw manure applications should occur at least 120 days before harvest). Decomposing plant materials will activate a diverse pool of microbes, including those that break down organic matter into plant-available nutrients as well as others that compete with plant pathogens on the root surface.

Rotating between crop families can help prevent the buildup of diseases that overwinter in the soil. Rotation with a grain crop, preferably a sod that will be in place for one or more seasons, deprives disease-causing organisms of a host, and also contributes to a healthy soil structure that promotes vigorous plant growth. The same practices are effective for preventing the buildup of root damaging nematodes in the soil, but keep in mind that certain grain crops are also hosts for some nematode species. Rotating between crops with late and early season planting dates can help prevent the buildup of weed populations. Organic growers must attend to the connection between soil, nutrients, pests, and weeds to succeed. An excellent resource for additional information on soils and soil health is [Building Soils for Better Crops](#) by Fred Magdoff and Harold Van Es, 2010 (Link 10). For additional

information, refer to the [Cornell Soil Health](#) website (Link 11).

### 3. COVER CROPS

Unlike cash crops, which are grown for immediate economic benefit, cover crops are grown for their valuable effect on soil properties and on subsequent cash crops. Cover crops help maintain soil organic matter, improve soil tilth, prevent erosion and assist in nutrient management. They can also contribute to weed management, increase water infiltration, maintain populations of beneficial fungi, and may help control insects, diseases and nematodes. To be effective, cover crops should be treated as any other valuable crop on the farm, with their cultural requirements carefully considered including their cultural requirements, life span, mowing recommendations, incorporation methods, and susceptibility, tolerance, or antagonism to root pathogens and other pests. Some cover crops and cash crops share susceptibility to certain pathogens and nematodes. Careful planning and monitoring is required when choosing a cover crop sequence to avoid increasing pest problems in subsequent cash crops. See Tables 3.1 for more information on specific cover crops and Section 8: *Crop and Soil Nutrient Management* for more information about how cover crops fit into a nutrient management plan.

A certified organic farmer is required to plant certified organic cover crop seed. If, after contacting at least three suppliers, organic seed is not available, then the certifier may allow conventional seed to be used. Suppliers should provide a purity test for cover crop seed. Always inspect the seed for contamination with weed seeds and return if it is not clean. Cover crop seed is a common route for introduction of new weed species onto farms.

#### 3.1 Goals and Timing for Cover Crops

Adding cover crops regularly to the crop rotation plan can result in increased yields of the subsequent cash crop. Goals should be established for choosing a cover crop; for example, the crop can add nitrogen, smother weeds, or break a pest cycle. The cover crop might best achieve some of these goals if it is in place for the entire growing season. If this is impractical, a compromise might be to grow the cover crop between summer cash crops. Allow two or more weeks between cover crop incorporation and cash crop seeding to permit decomposition of the cover crop, which will improve the seedbed and help avoid any unwanted allelopathic effects on the next cash crop. Another option is to overlap the cover crop and the cash crop life cycles by overseeding, interseeding or intercropping the cover crop between cash crop rows at final cultivation. An excellent resource for determining the best cover crop for your situation is

[Northeast Cover Crop Handbook](#), by Marianne Sarrantonio (Reference 6) or the [Cornell online decision tool](#) to match goals, season, and cover crop (Link 9).

Leaving cover crop residue to remain on the soil surface might make it easier to fit into a crop rotation and will help to conserve soil moisture, but some of the nitrogen contained in the residue will be lost to the atmosphere, and total organic matter added to the soil will be reduced. Turning under the cover crop will speed up the decomposition and nitrogen release from the crop residue.

#### 3.2 Legumes Cover Crops

Legume cover crops should be avoided before beans because many are closely related to beans and share pests.

#### 3.3 Non-Legume Cover Crops

Barley, rye grain, rye grass, Sudangrass, wheat, oats, and other grain crops left on the surface or plowed under as green manures or dry residue in the spring are beneficial because these plants take up nitrogen that otherwise might be leached from the soil, and release it back to the soil as they decompose. If incorporated, allow two weeks or more for decomposition prior to planting to avoid the negative impact on stand establishment from actively decomposing material. Three weeks might not be enough if soils are very cold. In wet years, the presence of cover crop residues may increase slug damage and infections by fungal pathogens such as *Pythium* and *Rhizoctonia*, affecting stand establishment

#### 3.4 Biofumigant Cover Crops

Certain cover crops have been shown to inhibit weeds, pathogens, and nematodes by releasing toxic volatile chemicals when tilled into the soil as green manures and degraded by microbes or when cells are broken down by finely chopping. Degradation is quickest when soil is warm and moist. These biofumigant cover crops include Sudangrass, sorghum-sudangrasses, and many in the brassica family. Varieties of mustard and arugula developed with high glucosinolate levels that maximize biofumigant activity have been commercialized (e.g. Caliente brands 199 and Nemat).

Attend to the cultural requirements of the cover crops to maximize growth. Fertilizer applied to the cover crops will be taken up and then returned to the soil for use by the cash crop after the cover crop is incorporated. Biofumigant cover crops like mustard should be allowed to grow to their full size, normally several weeks after flowering starts, but incorporated before the seeds become brown and hard indicating they are mature. To minimize loss of biofumigant, finely chop the tissue early in the day when temperatures are low. Incorporate immediately by tilling, preferably with a second tractor following the chopper. Lightly seal the soil

surface using a culti-packer and/or ½ inch of irrigation or rain water to help trap the volatiles and prolong their persistence in the soil. Wait at least two weeks before planting a subsequent crop to reduce the potential for the breakdown products to harm the crop, also known as phytotoxicity. Scratching the soil surface before planting will release remaining biofumigant. This biofumigant effect is not predictable or consistent. The levels of the active compounds and suppressiveness can vary by season, cover crop variety,

maturity at incorporation, amount of biomass, fineness of chopping, how quickly the tissue is incorporated, soil microbial diversity, soil tilth, and microbe population density.

[Cover Crops for Vegetable Growers: Decision Tool](#) (Link 9).

[Northeast Cover Crops Handbook](#) (Reference 6).

[Cover Crops for Vegetable Production in the Northeast](#) (Reference 11).

[Crop Rotation on Organic Farms: A Planning Manual](#) (Link 11a).

**Table 3.1. Non-Leguminous Cover Crops: Cultural Requirements and Crop Benefits**

SPECIES	PLANTING DATES	LIFE CYCLE	COLD HARDINESS ZONE (LINK 1)	HEAT	DROUGHT	SHADE	PH PREFERENCE	SOIL TYPE PREFERENCE	SEEDING (Lb/A)	COMMENTS
				--TOLERANCES--						
Brassicas e.g. mustards, rapeseed	April or late August-early Sept.	Annual / Biennial <sup>a</sup>	6-8	4	6	NI	5.3-6.8	Loam to clay	5-12	+Good dual purpose cover & forage +Establishes quickly in cool weather +Biofumigant properties
Buckwheat	Late spring-summer	Summer annual <sup>a</sup>	NFT	7-8	4	6	5.0-7.0	Most	35-134	+Rapid grower (warm season) +Good catch or smother crop +Good short-term soil improver for poor soils
Cereal Rye	August-early October	Winter annual	3	6	8	7	5.0-7.0	Sandy to clay loams	60-200	+Most cold-tolerant cover crop +Excellent allelopathic weed control +Good catch crop +Rapid germination & growth +Temporary N tie-up when turned under
Fine Fescues	Mid March-mid-May OR late Aug.-late Sept.	Long-lived perennial	4	3-5	7-9	7-8	5.3-7.5 (red) 5.0-6.0 (hard)	Most	16-100	+Very good low-maintenance permanent cover, especially in infertile, acid, droughty &/or shady sites
Oats	Mid-Sept-early October	Summer annual <sup>a</sup>	8	4	4	4	5.0-6.5	Silt & clay loams	110	+Rapid growth +Ideal quick cover and nurse crop
Ryegrasses	August-early Sept.	Winter annual (AR)/ Short-lived perennial (PR)	6 (AR) 4 (PR)	4	3	7 (AR) 5 (PR)	6.0-7.0	Most	14-35	+Temporary N tie-up when turned under +Rapid growth +Good catch crop +Heavy N & moisture users
Sorghum-Sudangrass	Late spring-summer	Summer Annual <sup>a</sup>	NFT	9	8	NI	Near neutral	NI	10-36	+Tremendous biomass producers in hot weather +Good catch or smother crop +Biofumigant properties

NI-No Information, NFT-No Frost Tolerance. Drought, Heat, Shade Tolerance Ratings: 1-2=low, 3-5=moderate, 6-8=high, 9-10=very high. . <sup>a</sup>Winter killed.

AR=Annual Rye, PR=Perennial Rye.

Reprinted with permission from [Rodale Institute](#) . M. Sarrantonio. 1994. Northeast Cover Crop Handbook. (Reference 6).

## 4. FIELD SELECTION

For organic production, give priority to fields with excellent soil tilth, high organic matter, good drainage and airflow. Beans do not thrive in wet soil.

### 4.1 Certifying Requirements

Certifying agencies have requirements that affect field selection. Fields cannot be treated with prohibited products for three years prior to the harvest of a certified organic crop. Adequate buffer zones are required between certified organic and conventionally grown crops. Buffer zones must be a barrier, such as a diversion ditch or dense hedgerow, or be a distance large enough to prevent drift of prohibited materials onto certified organic fields. Determining what buffer zone is needed will vary depending on equipment used on adjacent non-certified land. For example, use of high-pressure spray equipment or aerial pesticide applications in adjacent fields will increase the buffer zone size. Pollen from genetically engineered crops can also be a contaminant. An organic crop should not be grown near a genetically engineered crop of the same species. Check with your certifier for specific buffer requirements. These buffers commonly range between 20 to 250 feet depending on adjacent field practices.

### 4.2 Crop Rotation Plan

A careful crop rotation plan is the cornerstone of organic crop production because it allows the grower to improve soil quality and proactively manage pests. Although growing a wide range of crops complicates the crop rotation planning process, it ensures diversity in crop residues in the soil, and a greater variety of beneficial soil organisms. Individual organic farms vary widely in the crops grown and their ultimate goals, but some general rules apply to all organic farms regarding crop rotation. Rotating individual fields away from crops within the same family is critical and can help minimize crop-specific disease and non-mobile insect pests that persist in the soil or overwinter in the field or field borders. Pests that are persistent in the soil, have a wide host range, or are wind-borne, will be difficult to control through crop rotation. Conversely, the more host specific, non-mobile, and short-lived a pest is, the greater the ability to control it through crop rotation. The amount of time required for a crop rotation is based on the particular pest and its severity. Some particularly difficult pests may require a period of fallow. See specific recommendations in the disease and insect sections of this guide (Sections 11, 12, 13). Partitioning the farm into management units will help to organize crop rotations and ensure that all parts of the farm have sufficient breaks from each type of crop.

A well-planned crop rotation is key to weed management. Short season crops such as lettuce and spinach are harvested before many weeds go to seed, whereas vining cucurbits, with their limited cultivation time and long growing season, allow weeds to go to seed before harvest. Including short season crops in the rotation will help to reduce weed populations provided the field is cleaned up promptly after harvest. Other weed reducing rotation strategies include growing mulched crops, competitive cash crops, short-lived cover crops, or crops that can be intensively cultivated. Individual weed species emerge and mature at different times of the year, therefore alternating between spring, summer, and fall planted crops helps to interrupt weed life cycles.

Cash and cover crop sequences should also take into account the nutrient needs of different crops and the response of weeds to high nutrient levels. High soil phosphorus and potassium levels can exacerbate problem weed species. A cropping sequence that alternates crops with high and low nutrient requirements can help keep nutrients in balance. The crop with low nutrient requirements can help use up nutrients from a previous heavy feeder. A fall planting of a non-legume cover crop will help hold nitrogen not used by the previous crop. This nitrogen is then released when the cover crop is incorporated in the spring. See Section 5: *Weed Management*, and Section 3: *Cover Crops* for more specifics.

Rotating crops that produce abundant organic matter, such as hay crop and grain-legume cover crops, with ones that produce less, such as vegetables, will help to sustain organic matter levels and promote good soil tilth (see Section 2: *Soil Health* and Section 8: *Crop and Soil Nutrient Management*). Beans generally have a lower nutrient requirement (Table 4.2.1). Growing a cover crop, preferably one that includes a legume (unless the field has a history of *Pythium* or *Rhizoctonia* problems), prior to or after a bean crop, will help to renew soil nutrients, improve soil structure, and diversify soil organisms. Deep-rooted crops in the rotation to help break up compacted soil layers.

**Table 4.2.1 Crops Nutrient Requirements**

Crop	Nutrient Needs		
	Lower	Medium	Higher
	bean	cucumber	broccoli
	beet	eggplant	cabbage
	carrot	brassica greens	cauliflower
	herbs	pepper	corn
	pea	pumpkin	lettuce
	radish	spinach	potato
		chard	tomato
		squash	
		winter squash	

From NRAES publication [Crop Rotation on Organic Farms: A Planning Manual](#). Charles L. Mohler and Sue Ellen Johnson, editors, (Link 11a).

## Crop information specific to Beans

Beans should only be grown in fields that have had no legumes for 3 years. Legumes including soybean, clovers, alfalfa and hairy vetch are hosts for many soil-borne fungal pathogens and should be avoided in fields with severe root

rot problems. A good rotation helps reduce the incidence of foliar diseases and lowers the population of plant pathogens that cause root rot. Corn and cereal grains are excellent rotation crops to reduce root rot problems because they are not hosts for root rot pathogens of vegetable crops.

**Table 4.2.2 Potential Interactions of Crops Grown in Rotation with Beans**

Crops in Rotation	Potential Rotation Effects	Comments
Lettuce, potato, tomato, other legumes, crucifers, or cucurbits	Increase Sclerotinia	Avoid growing these crops prior to or after snap beans to reduce the buildup of Sclerotinia in the soil. Grow grains or corn for several years to help reduce <i>Sclerotinia</i> .
Bean, tomato, cucumber, buckwheat, aromatic herbs	Decrease Clubroot	Clubroot in brassicas has declined more quickly in fields where tomato, cucumber, snap bean and buckwheat have been grown. Aromatic perennial herbs such as summer savory, peppermint, or garden thyme helps to reduce clubroot when grown for 2 to 3 consecutive years.
Soybean, dry bean	Increases Bacterial blight ( <i>Xanthomonas campestris</i> )	<i>Xanthomonas campestris</i> is found in legumes as well as some crucifers and weeds in the mustard family.
Soybean, dry bean	Increases Soybean cyst nematode	Soybean cyst nematode <i>Heterodera glycines</i> increases to high densities on snap beans through the snap bean crop is rarely affected.
Snap bean, soybean, dry bean, alfalfa	Increase Fusarium, Pythium, and Sclerotinia	Do not plant these legumes prior to or for several years after snap beans to prevent a buildup of these soilborne diseases.

Excerpt from Appendix 2 of [Crop Rotation on Organic Farms: A Planning Manual](#), Charles L. Mohler and Sue Ellen Johnson, editors. (Link 11a)

## 4.3 Pest History

Knowledge about the pest history for each field is important for planning a successful cropping strategy. Germination may be reduced in fields with a history of *Pythium* or *Rhizoctonia*. Avoid fields that contain heavy infestations of perennial weeds such as nutsedge, bindweed, and quackgrass as these weeds are particularly difficult to control. One or more years focusing on weed population reduction using cultivated fallow and cover cropping may be needed before organic crops can be successfully grown in those fields. Susceptible crops should not be grown in fields with a history of *Sclerotinia* white mold without a rotation of several years to sweet corn or grain crops. If there is a field history of white mold, beans should not be preceded by tomato, potato, lettuce, crucifer crops, or bean (including soybean). Treat with Contans™ to reduce fungal sclerotia in the soil immediately after an infected crop is harvested.

If possible, beans should not be grown in fields with a history of root rot problems, but if there is no choice, plant as late as possible when the soil has warmed.

Root maggots prefer to lay eggs in soil with fresh organic matter. Incorporate cover crop residues 2-3 weeks before planting to allow time for decomposition.

Snap bean is a host for root-lesion nematode, *Pratylenchus penetrans*, and therefore it is important to know whether or not lesion nematode is present in the field in order to develop long-term crop rotations and cropping sequences that either reduce the populations in heavily infested fields or minimize their increase in fields that have no to low infestation levels. Refer to Section 12 for more information on nematodes.

## 4.4 Soil and Air Drainage

Most fungal and bacterial pathogens need free water on the plant tissue or high humidity for several hours in order to infect. Any practice that promotes leaf drying or drainage of excess water from the root zone will minimize favorable conditions for infection and disease development. Fields with poor air movement, such as those surrounded by hedgerows or woods, create an environment that favors prolonged leaf wetness. Plant rows parallel to the prevailing winds, which is typically in an east-west direction and avoid overcrowding by using wide row spacing to promote drying of the soil and reduce moisture in the plant canopy.

## 5. WEED MANAGEMENT

Weed management can be one of the biggest challenges on organic farms, especially during the transition and the first several years of organic production. To be successful,

weed management on organic farms must take an integrated approach that includes crop rotation, cover cropping, cultivation, and planting design, based on an understanding of dominant weed biology and ecology of dominant weed species. A multi-year approach that includes strategies for controlling problem weed species in a sequence of crops will generally be more successful than attempting to manage each year's weeds as they appear. Relying on cultivation alone to manage weeds in an organic system is a recipe for disaster.

Management plans should focus on the most challenging and potentially yield-limiting weed species in each field. Be sure, however, to emphasize options that do not increase other species that are present. Alternating between early and late-planted crops, and short and long season crops in the rotation can help minimize buildup of a particular weed or group of weeds with similar life cycles or growth habits, and will also provide windows for a variety of cover crops.

### 5.1 Record Keeping

Scout and develop a written inventory of weed species and their severity for each field. Accurate identification of weeds is essential. Weed fact sheets provide a good color reference for common weed identification. See Cornell [weed ecology](#) and Rutgers [weed gallery](#) websites (Link 13 and 14).

### 5.2 Weed Management Methods

Planting and cultivation equipment should be set up on the same number of rows to minimize crop losses and damage to crop roots during cultivation. It may be necessary to purchase specialized equipment to successfully control weeds in some crops.

Begin blind cultivation with a tine weeder, or flexible harrow, just before ground crack, when weeds are at white thread stage. Beans are very susceptible to breakage when they are in the "crook" stage – from just before ground crack until the seed leaves are unfolded and horizontal. Avoid tine weeding during this period. Correct cultivation depth is 2/3<sup>rd</sup> of seeding depth. Note, however, that penetration will vary with soil conditions and you must avoid hitting the seed with the weeder in soft spots. Effective tine weeding is an art that requires adjustment of the weeder to obtain good weed control without harming the crop. Examples of tine weeders are the Einbock, Lely and the Kovar. The tines on various brands and models of harrows differ in flexibility. Tines that are too stiff can break bean stems. Tines with a 70 to 80 degree bend work well for beans as they hook out grassy weeds without pulling out the beans, which have a taproot. Tines with a 45-degree bend can also be used effectively.

After bean emergence, make two more passes using a tine weeder at about 5 to 7 day intervals depending on weed

growth. Tine weeders work best on very small weeds. The final tine weeding can be more aggressive (faster and deeper) than the pre-emergence weeding or early post-emergence weeding. Test settings on a small area and adjust.

The flex tine harrow is effective with weeds in the row, but when crop reaches the early first trifoliate stage, the tine weeders may cause damage to the crop. At this point, a row crop cultivator can be used to control weeds between the rows. Adjust row crop cultivator for close and shallow cultivation. Cultivate to minimize disturbing the soil and uprooting rocks that will cause problems with the harvester. Perennial weeds will require deeper cultivation. Bean varieties that grow higher off the ground will minimize rocks in the harvester. Using a rolling cultivator is another option.

For a field with persistent perennial weeds such as field bindweed, several tactics might be needed to reduce competition: tillage to break up rhizomes, one or more short fallow periods to exhaust rhizome reserves, and planting a crop that requires multiple cultivations.

### Resources

[Steel in the Field](#) (Link 12).

[Cornell Weed Ecology website](#) (Link 13).

[Rutgers University, New Jersey Weed Gallery](#) (Link 14).

[Univ. of Vermont videos on cultivation and cover cropping](#) (Link 15).

[ATTRA Principles of Sustainable Weed Mgt. for Croplands](#) (Link 16).

[Cultivation Tools for Mechanical Weed Control in Vegetables](#) (Link 17)

## 6. RECOMMENDED VARIETIES

Variety selection is important both for the horticultural characteristics specified by the processor and the pest resistance profile that will be the foundation of a pest management program. Collaborate with processors on varieties, choosing those with some level of disease resistance if possible. Cornell research on developing CMV-resistant snap bean varieties is ongoing.

A certified organic farmer is required to plant certified organic seed. If, after contacting at least three suppliers, organic seed is not available for a particular variety, then the certifier may allow untreated conventional seed to be used.

**6.1.1 Varieties Currently Grown for Processing in NY**

Large Sieve Green	Whole Bean Type (2-3 sieve)
Huntington BA1001 Envy Venture	Flavor Sweet Masai Oakley
3-4 Sieve Green	Romano Type
Bowie Cabot Caprice Cassidy Colter	Furnano
	Wax Bean Type
	Gold Ribbon

**6.1.2 Varieties Identified as Candidates for Organic Production<sup>1</sup>**

3-4 Sieve Green	Whole Bean Type (2-3 sieve)
Bowie Caprice Cassidy Lewis	Banga Oakley

<sup>1</sup>Suggested for trial use only until more information is available

**7. PLANTING METHODS**

Recommended earliest planting date for untreated snap bean seed is June 1. The crop matures in 50 to 60 days, depending on the specific variety and desired pod size. Only western-grown, certified seed should be planted. Optimal germination of snap bean seed occurs at soil temperatures of 75° to 80°F. The minimum temperature at which snap bean germination will occur is 55° to 60°F. Plant rows in an east-west direction if possible and use wide row spacing, 36 inches, 5-7 plants/foot, to promote drying of the soil, increase air circulation, reduce moisture in the plant canopy and reduce risk of foliar diseases. Bean seed is sensitive to chilling during the initial stage of germination. If the soil is cold at this time, permanent damage may occur.

A good rotation helps reduce the incidence of foliar diseases and lowers the population of plant pathogens that cause root rot. Corn and cereal grains are excellent rotation crops. Planting on raised beds or ridges will help reduce root rot severity because the soil will be warmer and drier than the unridged soil. To allow for adequate aeration and drainage of excess moisture, avoid compacting the soil.

A water deficiency resulting from a lack of soil moisture or excessive transpiration can lead to deformed or pithy snap bean pods. Both yield and quality can be increased by irrigation before bloom and during pod enlargement if there is moisture stress. Irrigation during bloom with irrigation guns

that produce large droplets is not advised because blossoms can be knocked off the plant.

**8. CROP & SOIL NUTRIENT MANAGEMENT**

To produce a healthy crop, soluble nutrients must be available from the soil in amounts that meet the minimum requirements for the whole plant. The total nutrient needs of a crop are much higher than just the nutrients that are removed from the field when that crop is harvested. All of the roots, stems, leaves and other plant parts require nutrients at specific times during plant growth and development. The challenge in organic systems is balancing soil fertility to supply these required plant nutrients at a time and at sufficient levels to support healthy plant growth. Restrictions in any one of the needed nutrients will slow growth and can reduce crop quality and yields.

Organic growers often speak of feeding the soil rather than feeding the plant. A more accurate statement is that organic growers focus their fertility program on feeding soil microorganisms rather than the plant. Soil microbes decompose organic matter to release nutrients and convert organic matter to more stable forms such as humus. This breakdown of soil organic matter occurs throughout the growing season, depending on soil temperatures, water availability and soil quality. The released nutrients are then held on soil particles or humus making them available to crops or cover crops for plant growth. Amending soils with compost, cover crops, or crop residues also provides a food source for soil microorganisms and when turned into the soil, starts the nutrient cycle again.

During the transition years and the early years of organic production, soil amendment with composts or animal manure can be a productive strategy for building organic matter, biological activity and soil nutrient levels. This practice of heavy compost or manure use is not, however, sustainable in the long-term. If composts and manures are applied in the amounts required to meet the nitrogen needs of the crop, phosphorous may be added at higher levels than required by most vegetable crops. This excess phosphorous will gradually build up to excessive levels, increasing risks of water pollution or invigorating weeds like purslane and pigweed. A more sustainable, long-term approach is to rely more on legume cover crops to supply most of the nitrogen needed by the crop and use grain or grass cover crops to capture excess nitrogen released from organic matter at the end of the season to minimize nitrogen losses to leaching. See Section 3: *Cover Crops*. When these cover crops are incorporated into the soil, their nitrogen, as well as carbon, feeds soil microorganisms, supporting the nutrient cycle. Harvesting alfalfa hay from the field for several years can reduce high phosphorus and potassium levels.

The primary challenge in organic systems is synchronizing nutrient release from organic sources, particularly nitrogen, with the crop requirements. In cool soils, microorganisms are less active, and nutrient release may be too slow to meet the crop needs. Once the soil warms, nutrient release may exceed crop needs. In a long-term organic nutrient management approach, most of the required crop nutrients would be in place as organic matter before the growing season starts. Nutrients required by the crop in the early season can be supplemented by highly soluble organic amendments such as poultry manure composts or organically approved bagged fertilizer products (see Tables 8.2.4 to 8.2.6). These products can be expensive so are most efficiently used if banded at planting. The National Organic Standards Board states that no more than 20% of total N can be applied as Chilean nitrate. Be sure to confirm the practice with your organic certifier prior to field application.

Regular soil testing helps monitor nutrient levels, in particular phosphorus (P) and potassium (K). Choose a reputable soil-testing lab (Table 8.0.1) and use it consistently to avoid discrepancies caused by different soil extraction methods. Maintaining a soil pH between 6.3 and 6.8 will maximize the availability of all nutrients to plants.

**Table 8.0.1 Nutrient Testing Laboratories**

TESTING LABORATORY	SOIL	COMPOST/ MANURE	FORAGE	LINK
<a href="#">Cornell Soil Nutrient Analysis Lab</a>	x	x		18
<a href="#">Agri Analysis, Inc.</a>		x		19
<a href="#">A&amp;L Eastern Ag Laboratories, Inc.</a>	x	x		20
<a href="#">Penn State Ag Analytical Services Lab.</a>	x	x		21
<a href="#">The Agro-One Lab</a>			x	22
<a href="#">University of Massachusetts</a>	x	x		23

Develop a plan for estimating the amount of nutrients that will be released from soil organic matter, cover crops, compost, and manure. A strategy for doing this is outlined in Section 8.2: *Preparing an Organic Nutrient Budget*.

### 8.1 Fertility

Recommendations from the Cornell Integrated Crop and Pest Management Guidelines indicate that a bean crop requires 40 lbs. of available nitrogen (N), 80 lbs. of phosphorus (P), and 60 lbs. of potassium (K) per acre. These levels are based on the total needs of the whole plant and assume the use of synthetic fertilizers. Farmer and research experience suggests that lower levels may be adequate in organic systems. See Table 8.2.2 for the recommended rates of P and K based on soil test results. Nitrogen is not included because levels of available N change in response to soil

temperature and moisture, N mineralization potential, and leaching. As many of the nutrients as possible should come from cover crop, manure, and compost additions in previous seasons.

The source of these nutrients depends on soil type and historic soil management. Some soils are naturally high in P and K, or have a history of manure applications that have resulted in elevated levels. Additional plant available nutrients are supplied by decomposed soil organic matter or through specific soluble nutrient amendments applied during the growing season in organically managed systems. Many types of organic fertilizers are available to supplement the nutrients supplied by the soil. **ALWAYS check with your certifier before using any product to be sure it is approved.**

### 8.2 Preparing an Organic Nutrient Budget

Insuring an adequate supply of nutrients when the crop needs them requires careful planning. Developing an organic nitrogen budget can help estimate the amount of nutrients released by various organic amendments as well as native soil organic matter. Table 8.2.3 estimates common nutrient content in animal manures; however actual compost and manure nutrient content should be tested just prior to application. Analysis of other amendments, as well as cover crops, can be estimated using published values (see Tables 8.2.4 to 8.2.6 and 3.1 for examples). Keeping records of these nutrient inputs and subsequent crop performance will help evaluate if the plan is providing adequate fertility during the season to meet production goals.

Remember that with a long-term approach to organic soil fertility, the N mineralization rates of the soil will increase. This means that more N will be available from organic amendments because of increased soil microbial activity and diversity. Feeding these organisms different types of organic matter is essential to building this type of diverse biological community and ensuring long-term organic soil and crop productivity. Consider submitting soil samples for a [Cornell Soil Health Test](#) (Link 11). This test includes an estimate of nitrogen mineralization rate, which indicates the potential for release of N from soil organic matter. Testing soils over time can be useful for monitoring changes in nitrogen mineralization rate during the transition, and over time, in organic production.

Estimating total nutrient release from the soil and comparing it with soil test results and recommendations requires record-keeping and some simple calculations. Table 8.2.1 below can be used as a worksheet for calculating nutrients supplied by the soil compared to the total crop needs.

**Table 8.2.1 Calculating Nutrient Credits and Needs.**

	Nitrogen (N) lbs/A	Phosphate (P <sub>2</sub> O <sub>5</sub> ) lbs/A	Potash (K <sub>2</sub> O) lbs/A
1. Total crop nutrient needs			
2. Recommendations based on soil test	Not provided		
3. Credits			
a. Soil organic matter		---	---
b. Manure			
c. Compost			
d. Prior cover crop			
4. Total credits:			
5. Additional needs (2-4=)			

**Line 1. Total Crop Nutrient Needs:** Research indicates that an average bean crop requires 40 lbs. of available nitrogen (N), 80 lbs. of phosphorus (P), and 60 lbs. of potassium (K) per acre to support a medium to high yield crop (see Section 8.1 *Fertility* above).

**Line 2. Recommendations Based on Soil Test:** Use Table 8.2.2 to determine the amount of P and K needed based on soil test results.

**Table 8.2.2 Recommended Amounts of Phosphorus and Potassium for Beans Based on Soil Tests**

	Soil Phosphorus Level			Soil Potassium Level		
	low	med	high	low	med	high
Level shown in soil test						
	P <sub>2</sub> O <sub>5</sub> lbs/A			K <sub>2</sub> O lbs/A		
Total nutrient recommendation	80	60	40	60	40	20

**Line 3a. Soil Organic Matter:** Using the values from your soil test, estimate that 20 lbs. of nitrogen will be released from each percent organic matter in the soil. For example, a soil

that has 3% organic matter could be expected to provide 60 pounds of N per acre.

**Line 3b. Manure:** Assume that applied manure will release N for 3 years. Based on the test of total N in any manure applied, estimate that 50% is available in the first year, and then 50% of the remaining is released in each of the next two years. For an application rate of 100 lbs. of N as manure, 50 lbs. would be available the first year, 25 lbs. in year 2, and 12.5 lbs. in year 3. The NOP rules allow manure applications 120 days or more before harvest of a vegetable crop, but your farm certifier may have a more restrictive policy.

**Line 3c. Compost:** Estimate that between 10 and 25% of the N contained in compost will be available the first year. Compost maturity will influence how much N is available. If the material is immature, more of the N may be available to the crop in the first year. A word of caution: Using compost to provide for a crop's nutrient needs is not generally a financially viable strategy. The total volume needed can be very expensive for the units of N available to the crop, especially if trucking is required. Most stable composts should be considered as soil conditioners, improving soil health, microbial diversity, tilth, and nutrient retaining capacity. Any compost applied on organic farms must be approved for use by your farm certifier. Compost generated on the farm must follow an approved process outlined by your certifier.

**Line 3d. Cover Crops:** Legume cover crops are a source of nitrogen but are not recommended prior to a bean crop.

**Line 4. Total Credits:** Add together the various N values from the organic matter, compost, and cover crops to estimate the N supplying potential of the soil (see example below). There is no guarantee that these amounts will actually be available in the season, since soil temperatures, water, and crop physiology all impact the release and uptake of these soil nutrients.

Table 8.2.3 includes general estimates of nutrient availability for manures and composts but these can vary widely depending on animal feed, management of grazing, the age of the manure, amount and type of bedding, and many other factors. See table 3.1 for estimates of the nitrogen content of various cover crops. **Manure applications may not be allowed by your certifier or marketer even if applied 120 days before harvest. Check with both these sources prior to making manure applications.**

**Table 8.2.3 Nutrient Content of Common Animal Manures and Manure Composts**

	TOTAL N	P <sub>2</sub> O <sub>5</sub>	K <sub>2</sub> O	N1 <sup>1</sup>	N2 <sup>2</sup>	P <sub>2</sub> O <sub>5</sub>	K <sub>2</sub> O
	NUTRIENT CONTENT LB/TON			AVAILABLE NUTRIENTS LB/TON IN FIRST SEASON			
Dairy (with bedding)	9	4	10	6	2	3	9
Horse (with bedding)	14	4	14	6	3	3	13
Poultry (with litter)	56	45	34	45	16	36	31
Composted dairy manure	12	12	26	3	2	10	23
Composted poultry manure	17	39	23	6	5	31	21
Pelleted poultry manure <sup>3</sup>	80	104	48	40	40	83	43
Swine (no bedding)	10	9	8	8	3	7	7
	NUTRIENT CONTENT LB/1000 GAL.			AVAILABLE NUTRIENTS LB/1000 GAL FIRST SEASON			
Swine finishing (liquid)	50	55	25	25 <sup>4</sup>	20 <sup>5</sup>	44	23
Dairy (liquid)	28	13	25	14 <sup>4</sup>	11 <sup>5</sup>	10	23

1-N1 is an estimate of the total N available for plant uptake when manure is incorporated within 12 hours of application, 2-N2 is an estimate of the total N available for plant uptake when manure is incorporated after 7 days. 3 –Pelletized poultry manure compost. (Available in New York from Kreher's.)

4- injected, 5- incorporated.

Adapted from "Using Manure and Compost as Nutrient Sources for Fruit and Vegetable Crops" by Carl Rosen and Peter Bierman (Link 25) and [Penn State Agronomy Guide](#) 2013 (Link 25a).

Tables 8.2.4-8.2.6 lists some commonly available fertilizers and their nutrient content.

**Table 8.2.4 Available Nitrogen in Organic Fertilizer**

	Pounds of Fertilizer/Acre to Provide X Pounds of N per Acre				
Sources	20	40	60	80	100
Blood meal, 13% N	150	310	460	620	770
Soy meal 6% N (x 1.5) <sup>1</sup> also contains 2% P and 3% K <sub>2</sub> O	500	1000	1500	2000	2500
Fish meal 9% N, also contains 6% P <sub>2</sub> O <sub>5</sub>	220	440	670	890	1100
Alfalfa meal 2.5% N also contains 2% P and 2% K <sub>2</sub> O	800	1600	2400	3200	4000
Feather meal, 15% N (x 1.5) <sup>1</sup>	200	400	600	800	1000
Chilean nitrate 16% N cannot exceed 20% of crop's need.	125	250	375	500	625

<sup>1</sup> Application rates for some materials are multiplied to adjust for their slow to very slow release rates.

**Table 8.2.5 Available Phosphorous in Organic Fertilizer**

	Pounds of Fertilizer/Acre to Provide X Pounds of P <sub>2</sub> O <sub>5</sub> Per Acre				
SOURCES	20	40	60	80	100
Bonemeal 15% P <sub>2</sub> O <sub>5</sub>	130	270	400	530	670
Rock Phosphate 30% total P <sub>2</sub> O <sub>5</sub> (x4) <sup>1</sup>	270	530	800	1100	1300
Fish meal, 6% P <sub>2</sub> O <sub>5</sub> (also contains 9% N)	330	670	1000	1330	1670

<sup>1</sup> Application rates for some materials are multiplied to adjust for their slow to very slow release rates.

**Table 8.2.6 Available Potassium in Organic Fertilizers.**

	Pounds of Fertilizer/Acre to Provide X Pounds of K <sub>2</sub> O per acre:				
SOURCES	20	40	60	80	100
Sul-Po-Mag 22% K <sub>2</sub> O also contains 11% Mg	90	180	270	360	450
Wood ash (dry, fine, grey) 5% K <sub>2</sub> O, also raises pH	400	800	1200	1600	2000
Alfalfa meal 2% K <sub>2</sub> O also contains 2.5% N	1000	2000	3000	4000	5000
Greensand or Granite dust 1% K <sub>2</sub> O (x 4) <sup>1</sup>	8000	16000	24000	32000	40000
Potassium sulfate 50% K <sub>2</sub> O	40	80	120	160	200

<sup>1</sup> Application rates for some materials are multiplied to adjust for their slow to very slow release rates. Tables 8.4 to 8.6 adapted by Vern Grubinger from the [University of Maine soil testing lab](#) (Link 24).

### An example of how to determine nutrient needs for beans.

You will be growing an acre of snap beans. The Cornell Integrated Crop and Pest Management Guidelines suggests a total of 40 lb. N, 80 lb. P, and 60 lb. K per acre. Your soil tests show a pH of 6.5, with medium P and K levels and recommends 60 lbs P<sub>2</sub>O<sub>5</sub>/acre and 40 lbs K<sub>2</sub>O/acre (see Table 8.2.2). The field you will plant has 3% organic matter which will supply approximately 60 lbs N (line 3a). Last fall you spread but did not promptly incorporate 5 tons/acre of dairy manure with bedding before planting a rye cover crop

supplying ~10 lbs N (line 3d). Nutrient credits for soil organic matter, manure, and cover crop appear in Table 8.2.7.

**Table 8.2.7 Bean Example: Calculating Nutrient Credits and Needs Based on Soil Sample Recommendations.**

	Nitrogen (N) lbs/acre	Phosphate (P <sub>2</sub> O <sub>5</sub> ) lbs/acre	Potash (K <sub>2</sub> O) lbs/acre
1. Total crop nutrient needs:	40	80	60
2. Recommendations based on soil test	Not provided	60	40
3. Credits			
a. Soil organic matter 3%	60	-	-
b. Manure – 5 ton dairy	10	15	45
c. Compost - none	0	0	0
d. Cover crop – rye	0	0	0
4. Total credits:	70	15	45
5. Additional needed (2-4) =	0	45	0

Because of the threat of seedcorn maggot and possible N tie-up from the decomposing rye cover, wait at least three weeks between plowing and planting. This would be a good opportunity to do a shallow tine weeding just before planting. The dairy manure and soil organic matter will release N and K at adequate levels in this example; P is the only nutrient that needs to be applied. Because beans are sometimes planted into cold soils consider applying the 45 lbs/acre of P in a relatively soluble form such as composted chicken manure, which contains about 100 lbs P<sub>2</sub>O<sub>5</sub> per ton. Thus, banding about 900 lb at planting (2-3 inches to the side and below the furrow) would provide the P needed in this example, but would provide 40 lb of additional N, which is already higher than needed. Banding about 600 lbs of rock phosphate instead could avoid too much N, but the P would be less available in colder soils.

#### Additional Resources

[Using Organic Nutrient Sources](#) (reference 25b)

[Determining Nutrient Applications for Organic Vegetables](#) (reference 25c)

## 9. HARVESTING

All processing snap bean acreage is harvested by machine. Snap beans are processed relatively soon after harvest.

## 10. USING ORGANIC PESTICIDES

Given the high cost of many pesticides and the limited amount of efficacy data from replicated trials with organic products, the importance of developing an effective system of

cultural practices for insect and disease management cannot be emphasized strongly enough. **Pesticides should not be relied on as a primary method of pest control.** Scouting and forecasting are important for detecting symptoms of diseases at an early stage. When conditions do warrant an application, proper choice of materials, proper timing, and excellent spray coverage are essential.

### 10.1 Sprayer Calibration and Application

Calibrating sprayers is especially critical when using organic pesticides since their effectiveness is sometimes limited. For this reason, they tend to require the best spraying conditions to be effective. Read the label carefully to be familiar with the unique requirements of some products, especially those with live biological organisms as their active ingredient (e.g. Contans WG). The active ingredients of some biological pesticides (e.g. Serenade) are actually metabolic byproducts of the organism. Calculating nozzle discharge and travel speed are two key components required for applying an accurate pesticide dose per acre. Applying too much pesticide is illegal, can be unsafe and is costly whereas applying too little can fail to control pests or lead to pesticide resistance.

#### Resources

[Cornell Integrated Crop and Pest Management Guidelines: Pesticide Information and Safety](#) (Link 56).

[Calibrating Backpack Sprayers](#) (Link 57).

[Agricultural Pocket Pesticide Calibration Guide](#) (Link 58).

[Knapsack Sprayers – General Guidelines for Use](#) (Link 59)

[Herbicide Application Using a Knapsack Sprayer](#) (Link 60) This publication is relevant for non-herbicide applications.

[Pesticide Environmental Stewardship Community Page \(reference 64\)](#)

[Pesticide Environmental Stewardship Website \(reference 65\)](#)

[Vegetable Spraying \(reference 66\)](#)

### 10.2 Regulatory Considerations

Organic production focuses on cultural, biological, and mechanical techniques to manage pests on the farm, but in some cases pesticides, which include repellents, allowed for organic production are needed. Pesticides mentioned in this organic production guide are registered by the United States Environmental Protection Agency (EPA) or meet the EPA requirements for a “minimum risk” pesticide. The pesticides mentioned in this guide are also registered by the New York State Department of Environmental Conservation (NYS DEC) for use in New York State. See Cornell’s [Product, Ingredient, and Manufacturer System website](#) (Link 2) for pesticides currently registered for use in NYS. Additional products may be available for use in other states.

To maintain organic certification, products applied must also comply with the National Organic Program (NOP) regulations

as set forth in [7 CFR Part 205, sections 600-606](#) (Link 63). The [Organic Materials Review Institute](#) (OMRI) (Link 3) is one organization that reviews products for compliance with the NOP regulations and publishes lists of compliant products, but other entities also make product assessments. Organic growers are not required to use only OMRI listed materials, but the list is a good starting point when searching for allowed pesticides.

Finally, farms grossing more than \$5,000 per year and labeling products as organic must be certified by a NOP accredited certifier who must approve any material applied for pest management. ALWAYS check with the certifier before applying any pest control products. Some certifiers will review products for NOP compliance.

Note that "home remedies" may not be used. Home remedies are products that may have properties that reduce the impact of pests. Examples of home remedies include the use of beer as bait to reduce slug damage in strawberries or dish detergent to reduce aphids on plants. These materials are not regulated as pesticides, are not exempt from registration, and are therefore not legal to use.

#### **Do you need to be a certified pesticide applicator?**

The Federal Insecticide, Fungicide, and Rodenticide Act (FIFRA) defines two categories of pesticides: general-use and restricted use. NYS DEC also defines additional restricted-use pesticides. Pesticide applicator certification is required to purchase and use restricted-use pesticides. Restricted-use pesticides mentioned in this guide are marked with an asterisk (\*). Farmers who purchase and use only general-use pesticides on property they own or rent do not need to be certified pesticide applicators. However, we do encourage anyone who applies pesticides to become certified.

**Worker Protection Standard training.** If the farm has employees who will be working in fields treated with a pesticide, they must be trained as workers or handlers as required by the federal Worker Protection Standard (WPS). Having a pesticide applicator certification is one of the qualifications needed to be a WPS trainer. Certified pesticide applicators meet the WPS training requirements. For more information on the Worker Protection Standard see: [How To Comply with the Worker Protection Standard](#) (Link 67). Find more information on pesticide applicator certification from the list of [State Pesticide Regulatory Agencies](#) (Link 68) or, in New York State, see the Cornell Pesticide Management Education Program website at <http://psep.cce.cornell.edu> (Link 69).

### **10.3 Optimizing Pesticide Effectiveness**

Information on the effectiveness of a particular pesticide against a given pest can sometimes be difficult to find. Some university researchers include pesticides approved for organic production in their trials; some manufacturers provide trial results on their web sites; some farmers have conducted trials on their own. Efficacy ratings for pesticides listed in this guide were summarized from university trials and are only provided for some products. The [Resource Guide for Organic Insect and Disease Management](#) (Reference 1) provides efficacy information for many approved materials.

In general, pesticides allowed for organic production may kill a smaller percentage of the pest population, could have a shorter residual, and may be quickly broken down in the environment. Read the pesticide label carefully to determine if water pH or hardness will negatively impact the pesticide's effectiveness. Use of a surfactant may improve organic pesticide performance. [Omri lists adjuvants](#) on their website under *Crop Management Tools and Production Aids* (Link 3). Regular scouting and accurate pest identification are essential for effective pest management. Thresholds used for conventional production may not be useful for organic systems because of the typically lower percent mortality and shorter residual of pesticides allowed for organic production. When pesticides are needed, it is important to target the most vulnerable stages of the pest. Thoroughly cover plant surfaces, especially in the case of insecticides, since many must be ingested to be effective. The use of pheromone traps or other monitoring or prediction techniques can provide an early warning for pest problems, and help effectively focus scouting efforts.

## **11. DISEASE MANAGEMENT**

In organic systems, cultural practices form the basis of a disease management program. Promote plant health by maintaining a biologically active, well-structured, adequately drained and aerated soil that supplies the requisite amount and balance of nutrients. Choose varieties resistant to one or more important diseases whenever possible. Plant only clean, disease-free seed and maintain the best growing conditions possible.

Rotation is an important management practice for pathogens that overwinter in crop debris. Rotating between crop families is useful for many diseases, but may not be effective for pathogens with a wide host range, such as *Sclerotinia* white mold. Rotation with a grain crop, preferably a sod that will be in place for one or more seasons, deprives disease-causing organisms of a host, and also contributes to a healthy soil structure that promotes vigorous plant growth. The same

practices are effective for preventing the buildup of root damaging nematodes in the soil, but keep in mind that certain grain crops are also hosts for some nematode species. See more on crop rotation in Section 4.2: *Crop Rotation Plan*.

Other important cultural practices can be found under each individual disease listed below. Maximizing air movement and leaf drying is a common theme. Many plant diseases are favored by long periods of leaf wetness. Any practice that promotes faster leaf drying, such as orienting rows with the prevailing wind, or using a wider row or plant spacing, can slow disease development. Fields surrounded by trees or brush that tend to hold moisture after rain, fog or dew, should be avoided if possible.

Scouting fields weekly is key to early detection and evaluating control measures. The earlier a disease is detected, the more likely it can be suppressed with organic fungicides. When available, scouting protocols can be found in the sections listed below for each individual disease. While following a systematic scouting plan, keep watch for other disease problems. Removing infected plants during scouting is possible on a small operation. Accurate identification of disease problems, especially recognizing whether they are caused by a bacterium or fungus, is essential for choosing an effective control strategy. Anticipate which diseases are likely to be problems that could affect yield and be ready to take control action as soon as symptoms are seen. Allowing pathogen populations to build can quickly lead to a situation where there are few or no options for control.

All currently available fungicides allowed for organic production are protectants meaning they must be present on the plant surface before disease inoculum arrives to effectively prevent infection. They have no activity on pathogens once they are inside the plant. A few fungicides induce plant resistance and must be applied several days in advance of infection to be effective. Biological products must be handled carefully to keep the microbes alive. Follow label instructions carefully to achieve the best results.

Contact your local cooperative extension office to see if newsletters and pest management updates are available for your region. For example, in western New York, the [Cornell Vegetable Program](#) offers subscriptions to *VegEdge* a report that gives timely information regarding crop development, pest activity and control. Enrollment in the [Eastern New York Commercial Horticulture Program](#) includes a subscription to *Produce Pages* and weekly seasonal newsletters for vegetables, tree fruit, grapes and small fruit. On Long Island, see the *Long Island Fruit and Vegetable Update*.

Organic farms must comply with all other regulations regarding pesticide applications. See Section 10. Using Organic Pesticides for details. **ALWAYS check with your organic farm certifier when planning pesticide applications.**

#### Resources:

[Cornell Vegetable MD Online](#) (Reference 7)

[Resource Guide for Organic Insect and Disease Management](#) (Reference 1)

At the time this guide was produced, the following materials were available in New York State for managing this pest and were allowable for organic production. Listing a pest on a pesticide label does not assure the pesticide's effectiveness. The registration status of pesticides can and does change. Pesticides must be currently registered with the New York State Department of Environmental Conservation (DEC) to be used legally in NY. However, pesticides meeting the federal requirements for minimum-risk (25(b)) pesticides do not require registration. Current NY pesticide registrations can be checked on the Pesticide Product, Ingredient, and Manufacturer System [\(PIMS\) website](#) (Link 2). ALWAYS CHECK WITH YOUR CERTIFIER before using a new product.

**Table 11.0 Pesticides Labeled for Organic Bean Disease Control**

Class of Compound Product name - <i>Active ingredient</i>	Bacterial Diseases			Bean Rust	Botrytis Grey Mold	Pod-Flecking Complex	Root Rot & Damping Off	Sclerotinia White Mold
	Bacterial Brown Spot	Common Bacterial Blight	Halo blight					
BIOLOGICALS								
Actinovate AG ( <i>Streptomyces lydicus</i> )					X	X	X	X
Actinovate STP Fungicide ( <i>Streptomyces lydicus</i> )							X	
BIO-TAM ( <i>Trichoderma asperellum</i> , <i>Trichoderma gamsii</i> )							X	
Contans WG ( <i>Coniothyrium minitans</i> )								X
Double Nickel 55 Biofungicide ( <i>Bacillus amyloliquefaciens</i> str. D747)				X	X		X	X
Double Nickel LC Biofungicide ( <i>Bacillus amyloliquefaciens</i> str. D747)				X	X		X	X

ORGANIC SNAP BEAN PRODUCTION

Class of Compound Product name - Active ingredient	Bacterial Diseases			Bean Rust	Botrytis Grey Mold	Pod-Flecking Complex	Root Rot & Damping Off	Sclerotinia White Mold
	Bacterial Brown Spot	Common Bacterial Blight	Halo blight					
Mycostop Biofungicide ( <i>Streptomyces griseoviridis</i> )							X	
Mycostop Mix ( <i>Streptomyces griseoviridis</i> )							X	
Optiva ( <i>Bacillus subtilis</i> str. QST 713)					X			X
Prestop Biofungicide Powder ( <i>Gliocladium catenulatum</i> str. J1446)					X	X	X	
Regalia Biofungicide ( <i>Reynoutria sachalinensis</i> )		X		X	X		X	X
RootShield® Granules ( <i>Trichoderma</i> )							X	
RootShield PLUS+ Granules ( <i>Trichoderma</i> )							X	
RootShield PLUS+ WP ( <i>Trichoderma</i> )							X	
Serenade ASO ( <i>Bacillus subtilis</i> )				X				X
Serenade MAX ( <i>Bacillus subtilis</i> )				X				X
Serenade Optimum ( <i>Bacillus subtilis</i> )					X			X
Serenade Soil ( <i>Bacillus subtilis</i> )							X	
SoilGard ( <i>Gliocladium virens</i> str. GL-21)							X	
Taegro ( <i>Bacillus subtilis</i> )							X	
Zonix ( <i>Rhamnolipid</i> Biosurfactant)							X	
<b>BOTANICALS</b>								
Cinnerate (cinnamon oil)				X				
Trilogy (neem extract)				X	X	X		X
<b>COPPER</b>								
Badge X2 (copper oxychloride, copper hydroxide)	X	X	X					
Basic Copper 53 (basic copper sulfate)	X	X	X					
Champ WG (copper hydroxide)	X	X	X					
CS 2005 (copper sulfate pentahydrate)	X	X	X					
Cueva Fungicide Concentrate (copper octanoate)	X	X	X		X			X
NuCop 50DF (copper hydroxide)	X	X	X					
Nu-Cop HB (copper hydroxide)	X	X	X					
Nu-Cop 50 WP (copper hydroxide)	X	X	X					
Nordox 75 WG (cuprous oxide)			X					
<b>SULFUR</b>								
Micro Sulf (sulfur)	X			X				
Microthiol Disperss (sulfur)	X			X				
<b>OIL</b>								
JMS Stylet-Oil (paraffinic oil)				X				
Organic JMS Stylet Oil (paraffinic oil)				X				
PureSpray Green (petroleum oil)				X				
<b>OTHER</b>								
EcoMate ARMICARB 0 (potassium bicarbonate)					X			
Heads Up Plant Protectant ( <i>Chenopodium quinoa</i> )							X	
OxiDate 2.0 (hydrogen dioxide, peroxyacetic acid)				X	X		X	X
PERpose Plus (hydrogen peroxide/dioxide)	X	X	X	X	X	X	X	X
TerraClean 5.0 (hydrogen dioxide, peroxyacetic acid)					X		X	

X- labeled in New York state and OMRI listed product

## 11.1 Sclerotinia White Mold, *Sclerotinia sclerotiorum*

**Time for concern:** From open blossoms through the end of harvest

**Key characteristics:** The fungus will initially attack bean blossoms, as blossoms are a readily available source of food. Symptoms appear as white, fluffy cottony growth on blossoms, stems and pods. As the fungus grows, mounds of white mycelium harden and darken. These dark, black structures become sclerotia that enable the fungus to overwinter. Bean blossoms are an excellent source of nutrients for the fungus. Therefore control measures must be initiated at bloom. See Link 26. See Cornell [photo](#), and [bulletin](#) (Links 27-28) and learn more at the [Dillard Lab Vegetable Pathology](#) website (Link 28a).

Management Option	Recommendation for White Mold
<b>Scouting/thresholds</b>	Scout field prior to harvest to determine the need for treatment with Contans WG after harvest to reduce overwintering inoculum. Keep an accurate history of white mold incidence and severity in all fields.
<b>Coverage</b>	The best coverage can be obtained by using a minimum of 50 gallons per acre and high pressure (100 to 200 psi). Thoroughly cover initials, buds, and blossoms.
<b>Resistant varieties</b>	No resistant varieties are available, however plant architecture influences disease development. Select varieties with open canopies that hold pods high off the ground.
<b>Crop rotation</b>	If there is a field history of white mold, beans should not be preceded by a bean, tomato, potato, lettuce, or crucifer crop for several years. Grains and corn are good rotation crops.
<b>Site selection</b>	Avoid planting in shaded areas and in small fields surrounded by trees; do not plant in fields that drain poorly or have a history of severe white mold.
<b>Planting</b>	Plant rows in an east-west direction and use wide row spacing, 36 inches, to promote drying of the soil and reduce moisture in the plant canopy.
<b>Fertilization</b>	Avoid over fertilization.
<b>Postharvest</b>	Incorporate crop debris immediately following harvest to allow soil microorganisms the opportunity to feed on the survival structures called sclerotia or degrade disease organisms/overwintering structures.
<b>Seed selection/treatment</b>	This is not currently a viable management options.
<b>Note(s)</b>	White mold tends to develop in dense plant canopies. The disease tends to be worse in fields where there is poor weed management, where leaves have mechanical damage or pesticide injury, and where dead leaves are on the ground. The fungus can grow on dead and living material. White mold tends to develop when wet weather is persistent.

## ORGANIC SNAP BEAN PRODUCTION

At the time this guide was produced, the following materials were available in New York State for managing this pest and were allowable for organic production. Listing a pest on a pesticide label does not assure the pesticide's effectiveness. The registration status of pesticides can and does change. Pesticides must be currently registered with the New York State Department of Environmental Conservation (DEC) to be used legally in NY. However, pesticides meeting the federal requirements for minimum-risk (25(b)) pesticides do not require registration. Current NY pesticide registrations can be checked on the Pesticide Product, Ingredient, and Manufacturer System ([PIMS website](#) (Link 2)). ALWAYS CHECK WITH YOUR CERTIFIER before using a new product.

**Table 11.1 Pesticides Labeled for Management of Sclerotinia White Mold**

Class of Compound Product (Active ingredient)	Rate/A	PHI (days)	REI (hours)	Efficacy	Comments
<b>BIOLOGICALS</b>					
Actinovate AG ( <i>Streptomyces lydicus</i> WYEC 108)	3-12 oz Foliar spray or soil drench	0	1hr or until spray is dry	3	<i>Streptomyces lydicus</i> products effective in 0/2 trials. Reapply every 7-14 days. Use a spreader-sticker.
Contans WG ( <i>Coniothyrium minitans</i> )	1-4 lbs Soil treatment	-	4	2	Effective in 4/11 trials. Apply Contans WG to a Sclerotinia-infected crop immediately following harvest at 1 lb/A and incorporate the debris into the soil or apply at 2 lb/acre to a planted crop right after planting followed by shallow incorporation (or irrigate) to about a 1 to 2 inch depth. Do not turn the soil profile after application of Contans WG to avoid bringing untreated soil containing viable sclerotia near the surface. The seller recommends applying Contans WG for at least 3 to 4 years to reduce soil levels, or every year a susceptible crop is grown in that field. Enhance storage life by keeping product in the refrigerator or freezer.
Double Nickel 55 Biofungicide ( <i>Bacillus amyloliquefaciens</i> str. D747)	0.125-1 lb Soil treatment 0.25-3 lb Foliar treatment	0	4	?	
Double Nickel LC Biofungicide ( <i>Bacillus amyloliquefaciens</i> str. D747)	0.5-4.5 pints Soil treatment 0.5-6 qts Foliar treatment	0	4	?	
Optiva ( <i>Bacillus subtilis</i> str. QST 713)	14-24 oz Foliar treatment	0	4	?	Repeat on 7-10 day intervals or as needed.
Regalia Biofungicide ( <i>Reynoutria sachalinensis</i> )	1 – 4 qts Foliar or in-furrow treatment	0	4	?	Apply every 7-14 days as needed. Foliar application.
Serenade ASO ( <i>Bacillus subtilis</i> )	2-6 quarts Foliar spray	0	4	3	Not effective in 1/1 trial.
Serenade MAX ( <i>Bacillus subtilis</i> )	1-3 lbs Foliar treatment	0	4	3	
Serenade Optimum ( <i>Bacillus subtilis</i> )	14-20 oz Foliar treatment	0	4	?	
<b>COPPER</b> Copper products effective in 1/2 trials.					
Cueva Fungicide Concentrate (copper octanoate)	See comments Foliar treatment	Up to day of harvest	4	2	Product is mixed at 0.5 to 2.0 gallons per 100 gallons of water and applied at 50 to 100 gallons of diluted spray

# ORGANIC SNAP BEAN PRODUCTION

**Table 11.1 Pesticides Labeled for Management of Sclerotinia White Mold**

Class of Compound Product (Active ingredient)	Rate/A	PHI (days)	REI (hours)	Efficacy	Comments
OTHER					
Oxidate 2.0 (hydrogen dioxide, peroxyacetic acid)	32 fl oz – 1 gal/100 gal water Foliar treatment	0	Until spray has dried	?	.
PERpose Plus (hydrogen peroxide/dioxide)	1 fl oz/gal (initial/curative) Foliar treatment 0.25-0.33 fl oz/gal (weekly preventative) Foliar treatment	-	Until Dry	3	Hydrogen peroxide products not effective in 1/1 trial. For initial or curative use, apply higher rate for 1 to 3 consecutive days. Then follow with weekly/preventative treatment.  For weekly or preventative treatments, apply lower rate every five to seven days. At first signs of disease, use curative rate then resume weekly preventative treatment.

Efficacy: 1- effective in half or more of recent university trials, 2- effective in less than half of recent university trials, 3-not effective in any known trials, ?- not reviewed or no research available. PHI = pre-harvest interval, REI = restricted-entry interval. - = pre-harvest interval isn't specified on label.

## 11.2 Botrytis Gray Mold, Botrytis cinerea

**Time for concern:** From open blossoms through the end of harvest

**Key characteristics:** Gray mold develops in dense plant canopies when the weather is warm and moist. Large necrotic lesions produce gray spores on the pods, leaves and stems. This disease is worse in fields where leaves have mechanical damage or other injury. See Cornell [photo](#) (Link 29) and learn more at the [Dillard Lab Vegetable Pathology](#) website (Link 28a).

Management Option	Recommendations for Botrytis Gray Mold
<b>Scouting/thresholds</b>	Botrytis cinerea can infect many species of plants. A reliable forecasting system has not been developed for gray mold on beans. However, here are a few helpful hints on the development of gray mold. Gray mold tends to develop in dense plant canopies. The disease tends to be worse in fields where leaves have mechanical damage and where dead leaves are on the ground. The fungus grows and produces spores on dead and living material. These spores will subsequently infect bean pods. Gray mold tends to develop when the weather is warm and moist. Begin scouting when the first buds are showing. Record the occurrence and severity of gray mold. See Reference 5.
<b>Coverage</b>	The best coverage can be obtained by using a minimum of 50 gallons per acre and high pressure (100 to 200 psi).
<b>Resistant varieties</b>	No resistant varieties are available.
<b>Crop rotation</b>	Regular crop rotation is recommended. Grains and corn are good rotation crops.
<b>Site selection</b>	Avoid planting in shaded areas and in small fields surrounded by trees; do not plant in fields that drain poorly.
<b>Planting</b>	Plant rows in an east-west direction, and use wide row spacing (36 inches) to promote drying of the soil and reduce moisture in the plant canopy.
<b>Fertilization</b>	Avoid over-fertilization with nitrogen.

# ORGANIC SNAP BEAN PRODUCTION

<b>Postharvest</b>	Incorporate debris immediately after harvest to hasten decomposition of the material.
<b>Seed selection/treatment</b>	This is not currently a viable management options.
<b>Note(s)</b>	Avoid mechanical damage to leaves.

At the time this guide was produced, the following materials were available in New York State for managing this pest and were allowable for organic production. Listing a pest on a pesticide label does not assure the pesticide's effectiveness. The registration status of pesticides can and does change. Pesticides must be currently registered with the New York State Department of Environmental Conservation (DEC) to be used legally in NY. However, pesticides meeting the federal requirements for minimum-risk (25(b)) pesticides do not require registration. Current NY pesticide registrations can be checked on the Pesticide Product, Ingredient, and Manufacturer System ([PIMS website](#) (Link 2). ALWAYS CHECK WITH YOUR CERTIFIER before using a new product.

<b>Table 11.2 Pesticides Labeled for Management of Botrytis Gray Mold</b>					
<b>Class of Compound Product (Active ingredient)</b>	<b>Rate/A</b>	<b>PHI (days)</b>	<b>REI (hours)</b>	<b>Efficacy</b>	<b>Comments</b>
<b>BIOLOGICALS</b>					
Actinovate AG ( <i>Streptomyces lydicus</i> )	3-12 oz Soil treatment	0	1 or when dry	?	
Double Nickel 55 Biofungicide ( <i>Bacillus amyloliquefaciens</i> str. D747)	0.25-3 lbs Foliar treatment	0	4	?	
Double Nickel LC Biofungicide ( <i>Bacillus amyloliquefaciens</i> str. D747)	0.5-6 qts Foliar treatment	0	4	?	
Optiva ( <i>Bacillus subtilis</i> str. QST 713)	14-24 oz Foliar treatment	0	4	?	Repeat on 7-10 day intervals or as needed.
Prestop Biofungicide Powder ( <i>Gliocladium catenulatum</i> str. J1446)	3.5 oz/5 gal water, 0.5 gal suspension per 100 sq ft. Foliar treatment	-	0	?	Apply only when no above-ground harvestable food commodities are present.
Regalia Biofungicide ( <i>Reynoutria sachalinensis</i> )	1 – 4 qts Foliar treatment	0	4	?	Apply every 7-14 days as needed.
Serenade Optimum ( <i>Bacillus subtilis</i> )	14-20 oz Foliar treatment	0	4	?	
<b>BOTANICALS</b>					
Trilogy (neem extract)	0.5-1.0% in 25-100 gal water/A Foliar treatment	Up to day of harvest	4	1	Effective in 1/1 trial. Apply sufficient water to achieve complete coverage of foliage. Repeat applications every 7-14 days. Maximum labeled use of 2 gal/acre/application.
<b>COPPER</b>					
Cueva Fungicide Concentrate (copper octanoate)	See comments	Up to day of harvest	4	?	Product is mixed at 0.5 to 2.0 gallons per 100 gallons of water and applied at 50 to 100 gallons of diluted spray mixture per acre.
<b>OTHER</b>					
EcoMate ARMICARB 0 (potassium bicarbonate)	2.5-5 lbs/100 gal water Foliar treatment	0	4	?	
Oxidate 2.0 (hydrogen dioxide, peroxyacetic acid)	32 fl oz – 1 gal/100 gal water. Foliar treatment	0	Until spray has dried	?	
PERpose Plus (hydrogen peroxide/dioxide)	1 fl oz/ gal(initial/curative) Foliar treatment	-	Until dry	3	Not effective in 1/1 trial.  For initial or curative use, apply higher rate for 1 to 3 consecutive days. Then follow with

# ORGANIC SNAP BEAN PRODUCTION

**Table 11.2 Pesticides Labeled for Management of Botrytis Gray Mold**

Class of Compound Product (Active ingredient)	Rate/A	PHI (days)	REI (hours)	Efficacy	Comments
	0.25 – 0.33 fl oz/ gal (weekly preventative) Foliar treatment				weekly/preventative treatment.  For weekly or preventative treatments, apply lower rate every five to seven days. At first signs of disease, use curative rate then resume weekly preventative treatment
TerraClean 5.0 <i>hydrogen dioxide, peroxyacetic acid</i>	128 fl. oz./100 gal water. Soil treatment prior to seeding or transplanting.  25 fl oz/200 gal water/1000 ft <sup>2</sup> soil treated. Soil drench with established plants or seedlings.	-	0	?	

Efficacy: 1- effective in half or more of recent university trials, 2- effective in less than half of recent university trials, 3-not effective in any known trials, ?- not reviewed or no research available. PHI = pre-harvest interval, REI = restricted-entry interval. - = pre-harvest interval isn't specified on label.

## 11.3 Pod-flecking complex (PFC), *Alternaria alternata* and *Plectosporium tabacinum*

**Time for concern:** From pod fill through the end of harvest

**Key characteristics:** Pod-flecking complex is sometimes referred to by consultants and producers as russet, seam rust, spots, or rusty or spotty beans. PFC symptoms on pods intensify with pod maturity, and are most prevalent mid- to late August following periods of prolonged rainfall or rainfall of high intensity. Symptoms on pods include tan, orange, or black discolorations in the suture and/or small dark superficial specks, flecks, or spots (sometimes sunken) on the pod surfaces. See more at the [Dillard Lab Vegetable Pathology](#) website (Link 28a).

Management Options	Recommendations for Pod-Flecking Complex
<b>Scouting/thresholds</b>	Pod-flecking complex can infect many species of plants. A reliable forecasting system has not been developed for PFC on beans. However, here are a few helpful hints on the development of PFC. Pod-flecking complex symptoms on pods intensify with pod maturity, and are most prevalent mid- to late August following periods of prolonged rainfall or rainfall of high intensity. Only the pods are affected. Symptoms include tan, orange, or black discolorations in the suture and/or small dark superficial specks, flecks, or spots (sometimes sunken) on the pod surfaces. Begin scouting at pod fill. Record the occurrence and severity of PFC. See Reference 12.
<b>Coverage</b>	The best coverage can be obtained by using a minimum of 50 gallons per acre and high pressure (100 to 200 psi).
<b>Resistant varieties</b>	No resistant varieties are available.
<b>Crop rotation</b>	Regular crop rotation is recommended to improve plant health.
<b>Site selection</b>	Avoid planting in shaded areas and in small fields surrounded by trees; do not plant in fields that drain poorly.
<b>Planting</b>	Plant rows in an east-west direction, and use wide row spacing (36 inches) to promote drying of the soil and reduce moisture in the plant canopy.

## ORGANIC SNAP BEAN PRODUCTION

<b>Fertilization</b>	Avoid over-fertilization with nitrogen.
<b>Harvest / Postharvest</b>	To mitigate disease, harvest at or near peak maturity and avoid harvest delays that would result in overripe pods. Don't store infected beans.  Incorporate debris immediately after harvest to hasten decomposition of the material.
<b>Seed selection/treatment</b>	This is not currently a viable management options.

At the time this guide was produced, the following materials were available in New York State for managing this pest and were allowable for organic production. Listing a pest on a pesticide label does not assure the pesticide's effectiveness. The registration status of pesticides can and does change. Pesticides must be currently registered with the New York State Department of Environmental Conservation (DEC) to be used legally in NY. However, pesticides meeting the federal requirements for minimum-risk (25(b)) pesticides do not require registration. Current NY pesticide registrations can be checked on the Pesticide Product, Ingredient, and Manufacturer System ([PIMS website](#)) (Link 2). ALWAYS CHECK WITH YOUR CERTIFIER before using a new product.

Table 11.3 Pesticides Labeled for Management of Pod-Flecking Complex					
Class of Compound Product (Active ingredient)	Rate/A	PHI (days)	REI (hours)	Efficacy	Comments
<b>BIOLOGICALS</b>					
Actinovate AG ( <i>Streptomyces lydicus</i> WYEC 108)	3-12 oz Foliar treatment	0	1 or until dry	?	Labeled for <i>Alternaria</i> only. Reapply every 7-14 days. The Label recommends use of a spreader-sticker for foliar sprays.
Prestop Biofungicide Powder ( <i>Gliocladium catenulatum</i> )	1.4-3.5 oz/2.5 gal water Soil drench.	-	0	?	Labeled for <i>Alternaria</i> only. Treat only the growth substrate when above-ground harvestable food commodities are present.
<b>BOTANICALS</b>					
Trilogy (neem extract)	0.5-1.0% in 25- 100 gal water/A Foliar treatment	Up to day of harvest	4	?	Maximum labeled use of 2 gal/A/application.

Efficacy: 1- effective in half or more of recent university trials, 2- effective in less than half of recent university trials, 3-not effective in any known trials, ?- not reviewed or no research available. PHI = pre-harvest interval, REI = restricted-entry interval. - = pre-harvest interval isn't specified on label.

### 11.4 Root Rot and Damping-Off.

*There are four pathogenic fungi, Pythium, Rhizoctonia, Thielaviopsis, and Fusarium that are often found attacking bean roots and causing yield losses.*

**Time for concern:** Planting through end of bloom stage. Long term planning is required for sustainable management.

**Key characteristics:** Pythium is seen early in the season when it causes rotting of seeds and damping-off of young seedlings. In older plants, it causes reduction and discoloration of the root system. Pythium can also attack pods, causing lesions that generally develop at the tip of the pod and expand toward the stem. See Cornell [pythium photo](#) (Link 31). Rhizoctonia is also capable of causing seed decay and damping-off diseases on seedlings. See Cornell [rhizoctonia photo](#) (Link 32). On older plants it produces reddish brown, sunken lesions on the stem and tap root, and is generally favored by warm soil conditions. In addition, the sexual stage of Rhizoctonia may also be detected on the stem, petioles, and pods as a thin, whitish, compact growth. Thielaviopsis is often referred to as black root rot because the initial elongated lesions and later large infected areas on the stems and roots are dark brown to charcoal. This disease is favored by hot, wet conditions. Fusarium rot causes longitudinal, brick red lesions on the stem and tap root and is very common. See Reference 9 and Cornell [Bulletin 110](#) (Link 30) for more photos and information.

# ORGANIC SNAP BEAN PRODUCTION

Management Option	Recommendations for Root Rot and Damping-Off
Scouting/thresholds	Record the occurrence, type, and severity of root rot. No thresholds are available.
Resistant varieties	All commercially acceptable varieties are susceptible, but a number yield better under severe root rot.
Crop rotation	Rotate away from vegetables. One or two years with a grain crop such as barley, oats, rye, wheat, or corn will prevent severe root rot development when practiced on relatively clean fields; longer rotations are necessary in heavily infested fields. Avoid planting legumes as rotational crops or cover crops in heavily infested areas.
Site selection	Choose healthy and well-drained soils. A soil-indexing procedure that differentiates relatively pathogen-free fields from those with severe root rot problems is available. See the Cornell <a href="#">Soil Health Assessment Manual</a> (Link 11) for directions to do this yourself or to send a sample to Cornell for testing.
Seed selection/treatment	Select vigorous, disease-free seeds.
Soil treatment	Breaking hard pans, plowing deep, or ripping and planting on raised ridges or beds will reduce damage from root rot diseases.
Cover crop	Barley, rye grain, rye grass, wheat, oats, and other grain crops left on surface or plowed under as green manures or dry residue in the spring are beneficial. If incorporated as green manures, allow 2 weeks or more for decomposition prior to planting. Sudangrass or sorghum sudangrass hybrids can also be used as green manures. In wet years, using green manures may increase slug damage and affect stand establishment.
Planting	Avoid planting in heavily infested fields, but if there is no choice, plant shallow and late. Plantings exhibiting symptoms of severe root rot damage will benefit from a shallow cultivation not too close to the stems. Also, covering the lower stem tissues with soil will promote further root formation and reduce root rot damage. However, the latter should be done on an emergency basis, as this practice has been observed to increase foliar infections with <i>Rhizoctonia</i> in wet seasons.
Postharvest	Crop debris should be plowed down to initiate decomposition, if tillage system in use permits.

At the time this guide was produced, the following materials were available in New York State for managing this pest and were allowable for organic production. Listing a pest on a pesticide label does not assure the pesticide's effectiveness. The registration status of pesticides can and does change. Pesticides must be currently registered with the New York State Department of Environmental Conservation (DEC) to be used legally in NY. However, pesticides meeting the federal requirements for minimum-risk (25(b)) pesticides do not require registration. Current NY pesticide registrations can be checked on the Pesticide Product, Ingredient, and Manufacturer System ([PIMS website](#) (Link 2)). ALWAYS CHECK WITH YOUR CERTIFIER before using a new product.

**Table 11.4 Pesticides Labeled for Management of Root Rot and Damping-Off**

Class of Compound Product (Active ingredient)	Rate/A	PHI (days)	REI (hours)	Efficacy	Comments
<b>BIOLOGICALS</b>					
Actinovate AG ( <i>Streptomyces lydicus</i> )	3-12 oz Soil treatment	0	1 or until dry	?	Not labeled for <i>Thielaviopsis</i> .
Actinovate STP Fungicide <i>Streptomyces lydicus</i>	4-32 oz/100 lb of seed Seed treatment	-	1 or until dry	?	
BIO-TAM ( <i>Trichoderma asperellum</i> , <i>Trichoderma gamsii</i> )	1.5-3 oz/1000 ft row In furrow application	-	1	?	

ORGANIC SNAP BEAN PRODUCTION

Table 11.4 Pesticides Labeled for Management of Root Rot and Damping-Off					
Class of Compound Product (Active ingredient)	Rate/A	PHI (days)	REI (hours)	Efficacy	Comments
Double Nickel 55 Biofungicide – ( <i>Bacillus amyloliquefaciens</i> str. D747)	0.125-1 lb Soil application	0	4	?	
Double Nickel LC Biofungicide – ( <i>Bacillus amyloliquefaciens</i> str. D747)	0.5-4.5 pints Soil application	0	4	?	
Mycostop Mix ( <i>Streptomyces griseoviridis</i> str. K61)	5-8 oz/100 lbs of seed Seed treatment  7.6-30 oz. Soil spray or drench  0.5-1 lb/ treated acre Band, in-furrow or side dress.	0	4	?	Labeled only for root rots and damping off caused by <i>Fusarium</i> spp.  Use at planting; no pre-harvest interval noted. Irrigate within 6 hours after soil spray or drench with enough water to move Mycostop into the root zone.  Lightly incorporate furrow or band applications.
Mycostop Biofungicide ( <i>Streptomyces griseoviridis</i> Strain K61)	7 oz/cwt seed Seed treatment  15-30 oz Soil spray or drench	-	4	?	Labeled only for root rots and damping off caused by <i>Fusarium</i> spp.  Irrigate within 6 hours after soil spray or drench with enough water to move Mycostop Biofungicide into the root zone.
Prestop Biofungicide Powder ( <i>Gliricladium catenulatum</i> str. J1446)	1.4-3.5 oz per 2.5 gal water Soil drench.	-	0	?	Treat only the growth substrate when above- ground harvestable food commodities are present. Not labeled for <i>Thielaviopsis</i> .
RootShield Granules ( <i>Trichoderma</i> )	5-12 lbs In-furrow application	-	0	2	<i>Trichoderma harzianum</i> products effective against seedling diseases in 1/8 trials. <i>T. virens</i> products effective against <i>Fusarium</i> root rot in 0/2 trials.
RootShield PLUS+ Granules Biological Fungicide ( <i>Trichoderma harzianum</i> and <i>T.</i> <i>virens</i> )	5-12 lbs/A In-furrow application.	-	0	2	<i>Trichoderma harzianum</i> products effective against seedling diseases in 1/8 trials. <i>T. virens</i> products effective against <i>Fusarium</i> root rot in 0/2 trials.
RootShield PLUS+ WP Biological Fungicide ( <i>Trichoderma harzianum</i> and <i>T.</i> <i>virens</i> ))	3-8 oz/100 gal water Chemigation application.	0	4	2	<i>Trichoderma harzianum</i> products effective against seedling diseases in 1/8 trials. <i>T. virens</i> products effective against <i>Fusarium</i> root rot in 0/2 trials. Do not apply when above-ground harvestable food commodities are present.
Regalia Biofungicide terracl ( <i>Reynoutria sachalinensis</i> )	1-4 qts In-furrow application.	0	4	?	Not labeled for <i>Thielaviopsis</i>
Serenade Soil ( <i>Bacillus subtilis</i> strain QST 713)	2-6 qts Soil surface drench.	0	4	?	Not labeled for <i>Thielaviopsis</i>
SoilGard Microbial Fungicide ( <i>Gliricladium virens</i> str. GL-21)	2-10 lbs/A in 50-100 gal water In-furrow application	-	0	?	
Taegro ( <i>Bacillus subtilis</i> var. <i>amyloliquefaciens</i> strain FZB24)	3 tsp/gallon of water Seed treatment  2.6 oz/ 100 gallons for 2 acres. Over furrow at time of planting	-	24	?	Only labeled for <i>Fusarium</i> and <i>Rhizoctonia</i> .

# ORGANIC SNAP BEAN PRODUCTION

**Table 11.4 Pesticides Labeled for Management of Root Rot and Damping-Off**

Class of Compound Product (Active ingredient)	Rate/A	PHI (days)	REI (hours)	Efficacy	Comments
Zonix (Rhamnolipid Biosurfactant)	0.5-0.8 oz/gal water Foliar treatment	-	4		Labeled for <i>Pythium</i> only.
<b>OTHER</b>					
Heads Up Plant Protectant ( <i>Chenopodium quinoa extract</i> )	.13 oz/gal water Seed treatment	-	12	?	Use 34 ounces of mixed product to treat 360 pounds of seed. Thoroughly coat seed surface with solution. Use immediately after mixing.
Oxdate 2.0 (hydrogen dioxide, peroxyacetic acid)	0.5-1 gal In-furrow application.	0	Until spray has dried	?	
PERpose Plus (hydrogen peroxide/dioxide)	1 fl oz/ gal (initial/curative) Soil drench  0.25 – 0.33 fl oz/ gal (weekly preventative) Soil drench  1 fl oz/gal water Soil drench	-	Until dry	?	For initial or curative use, apply higher rate for 1 to 3 consecutive days. Then follow with weekly/preventative treatment.  For weekly or preventative treatments, apply lower rate every five to seven days. At first signs of disease, use curative rate then resume weekly preventative treatment.
TerraClean 5.0 (hydrogen dioxide, peroxyacetic acid)	128 fl. oz./100 gal water Soil treatment prior to seeding or transplanting.  25 fl oz/200 gal water/1000 ft <sup>2</sup> soil treated. Soil drench with established plants or seedlings.	-	0	?	

Efficacy: 1- effective in half or more of recent university trials, 2- effective in less than half of recent university trials, 3-not effective in any known trials, ?- not reviewed or no research available PHI = pre-harvest interval, REI = restricted-entry interval. - = pre-harvest interval isn't specified on label.

## 11.5 Bacterial Diseases

**Bacterial brown spot** (*Pseudomonas syringae* pv. *syringae*); **Common bacterial blight**, (*Xanthomonas campestris* pv. *phaseoli*); and **Halo blight**, (*Pseudomonas syringae* pv. *phaseolicola*)

**Time for concern:** From seeding stage through the end of harvest

**Key characteristics:** **Brown spot** - small, brown spots, 3/25 to 8/25 inch in diameter, often with a narrow, diffuse, pale margin, appear on leaves. Sunken, brown spots can form on the pods. **Common bacterial blight** - light brown lesions of irregular shape with distinct, bright yellow margins, 2/5 inch long, appear on leaves. Spots form on the pods and enlarge into reddish brown lesions. In humid weather, yellow bacteria may be present on the lesions. **Halo blight** - small, water-soaked spots on the undersides of leaves develop into numerous, small, reddish brown lesions with pale to yellow margins or halos. Pod symptoms are similar to those of common blight. See Cornell [fact sheet](#) (Link 33).

Management Option	Recommendations for Bacterial Diseases
Scouting/thresholds	Fields should be scouted at least twice between midseason and harvest. Record the occurrence and severity of the bacterial blights.

# ORGANIC SNAP BEAN PRODUCTION

Management Option	Recommendations for Bacterial Diseases
<b>Resistant varieties</b>	Some varieties have tolerance and/or resistance to one or more of these diseases. Depending on the variety, brown spot development on leaves may or may not result in significant damage to pods. Wisconsin growers report that 'Hystyle' shows resistance to brown spot.
<b>Crop rotation</b>	In the case of halo blight, rotate away from fields where this disease has occurred for a minimum of three years. Use a two-year minimum rotation for bacterial brown spot or common bacterial blight. <i>Xanthomonas campestris</i> also harbors in weeds within the mustard family.
<b>Site selection</b>	Avoid planting snap bean fields near dry bean fields.
<b>Seed selection/treatment</b>	Plant only western-grown, certified seed.
<b>Postharvest</b>	Crop debris should be destroyed as soon as possible to remove this source of disease for future plantings and to initiate decomposition.
<b>Sanitation</b>	Equipment used in fields with bacterial diseases should be thoroughly cleaned before being moved to disease-free fields. To reduce the spread of bacteria on equipment or in spray water, avoid making pesticide applications or cultivating when the leaves are wet. If possible, plow under bean stubble immediately after harvest.

At the time this guide was produced, the following materials were available in New York State for managing this pest and were allowable for organic production. Listing a pest on a pesticide label does not assure the pesticide's effectiveness. The registration status of pesticides can and does change. Pesticides must be currently registered with the New York State Department of Environmental Conservation (DEC) to be used legally in NY. However, pesticides meeting the federal requirements for minimum-risk (25(b)) pesticides do not require registration. Current NY pesticide registrations can be checked on the Pesticide Product, Ingredient, and Manufacturer System ([PIMS website](#) (Link 2)). ALWAYS CHECK WITH YOUR CERTIFIER before using a new product.

**Table 11.5 Pesticides Labeled for Management of Bacterial Diseases**

Class of Compound Product (Active ingredient)	Rate/A	PHI (days)	REI (hours)	Efficacy	Comments
<b>BIOLOGICALS</b>					
Regalia Biofungicide ( <i>Reynoutria sachalinensis</i> )	1-4 qts Foliar treatment	0	4	?	Labeled only for <i>Xanthomonas</i> .
<b>COPPER COMPOUNDS</b>					
<b>Good control has been inconsistent with copper compounds.</b>					
Badge X2 (copper oxychloride, copper hydroxide)	0.5-1.25 lbs Foliar treatment	-	48	2	
Basic Copper 53 (basic copper sulfate)	2 – 4 lbs Foliar treatment	Up to day of harvest	24	2	If possible time applications so that 12 hours of dry weather follow application.
Champ WG (copper hydroxide)	1.58 lbs Foliar treatment	-	48	2	
CS 2005 (copper sulfate pentahydrate)	19.2 - 25.6 oz Foliar treatment	-	48	2	
Cueva Fungicide Concentrate (copper octanoate)	See comments	Up to day of harvest	4	2	Product is mixed at 0.5 to 2.0 gallons per 100 gallons of water and applied at 50 to 100 gallons of diluted spray mixture per acre.
Nordox 75 WG (cuprous oxide)	0.6 – 2.5 lbs Foliar treatment	-	12	2	Labeled only for halo blight. Apply when plants are 5-6 inches tall, repeat in 7-14 day intervals.
NuCop 50DF (copper hydroxide)	1-3 lbs Foliar treatment	1	24	2	

# ORGANIC SNAP BEAN PRODUCTION

Table 11.5 Pesticides Labeled for Management of Bacterial Diseases					
Class of Compound Product (Active ingredient)	Rate/A	PHI (days)	REI (hours)	Efficacy	Comments
Nu-Cop HB (copper hydroxide)	0.5-1.5 lbs Foliar treatment	1	24	?	Use higher rate for more severe disease. Repeat on 7-14 day schedule depending on local conditions.
Nu-Cop 50 WP (copper hydroxide)	1.5-3 lbs Foliar treatment	1	24	2	
<b>SULFUR</b>					
Micro Sulf (sulfur)	7 lbs Foliar treatment	-	24	?	Consult processor before using sulfur.
Microthiol Disperss (sulfur)	3-10 lbs Foliar treatment	-	24	?	Consult processor before using sulfur. Only labeled for bacterial brown spot.
<b>OTHER</b>					
PERpose Plus (hydrogen peroxide/dioxide)	1 fl oz/ gal(initial/curative) Foliar treatment  0.25 – 0.33 fl oz/ gal(weekly preventative) Foliar treatment	-	Until dry	?	For initial or curative use, apply higher rate for 1 to 3 consecutive days. Then follow with weekly/preventative treatment.  For weekly or preventative treatments, apply lower rate every five to seven days. At first signs of disease, use curative rate then resume weekly preventative treatment

Efficacy: 1- effective in half or more of recent university trials, 2- effective in less than half of recent university trials, 3-not effective in any known trials, ?- not reviewed or no research available PHI = pre-harvest interval, REI = restricted-entry interval. - = pre-harvest interval isn't specified on label.

## 11.6 Virus Diseases

Virus diseases of bean are spread by infected seed or by aphids. However, controlling aphids is not effective for reducing these viruses. Virus expression is lower when the crop has adequate water. Irrigating in dry years may help mitigate impacts of virus infection. Record the occurrence and severity of any viruses present. See Cornell [photos](#) (Link 35), [fact sheet](#) (Link 34), and a [list of weed and crop hosts](#) (Link 36). Information about aphid vector activity and spread of CMV in snap bean fields can be found in References 3 and 6. Cornell research on developing CMV-resistant snap bean varieties is on going.

Disease/Symptoms	Spread by	Time for concern	Resistant Varieties	Notes
Cucumber Mosaic Virus (CMV) Leaf curl, green mottle, blistering, and a rugged zipper-like appearance along the main veins involving only a few leaves. Infected plants may also not express symptoms.	Soybean aphid, yellow clover aphid, pea aphid, corn leaf aphid; rarely seed transmitted	Primarily in plantings from late June through late July.	None available	Use certified, disease-free seed. CMV does not persist in plant debris, in the soil, or on equipment. For CMV and the other aphid-transmitted viruses listed below, there will be less risk of yield loss if fields are planted early (mid May to late June) than if planted after this period.
Bean Common Mosaic Virus (BCMV) Symptoms include a green mosaic and downward cupping along the main vein of each leaflet. Green vein banding, blistering, and malformation are common in leaves of the same plant. Plants are small, and pods	Seed, bean aphid, cowpea aphid, pea aphid, potato aphid, green peach	Seeding through harvest	Most varieties carry the I-gene for resistance to BCMV.	Use certified, disease-free seed.

## ORGANIC SNAP BEAN PRODUCTION

Disease/Symptoms	Spread by	Time for concern	Resistant Varieties	Notes
may be mottled and malformed. Symptoms are persistent.	aphid			
Bean Yellow Mosaic Virus (BYMV) Leaf mosaic formed by contrasting yellow or green mosaic areas. Pods generally are not affected, but the number of seeds per pod may be reduced.	Bean aphid, cowpea aphid, pea aphid, potato aphid, and green peach aphid.	June through harvest	None available	BYMV does not persist in plant debris, in the soil, or on equipment.
Clover Yellow Vein Virus (CYVV) In addition to deforming pods, this virus also causes a prominent yellow mosaic, malformation, and reduction in plant size.	Aphids	June through harvest	None available	

### 11.7 Bean Rust, *Uromyces appendiculatus*

**Time for concern:** Early in the growing season

**Key characteristics:** Regular occurrence of dew favors infection and development of severe epidemics. Bean rust is characterized by reddish brown, circular pustules on leaves or pods. See University of Connecticut [fact sheet](#) (Link 37). Bean rust is rarely seen in New York. However, growers commonly use the name “rust” to describe a diffuse light brown discoloration that sometimes occurs on the pods. See section 10.3 on Pod-flecking complex (Reference 12). Also see the [Dillard Lab Vegetable Pathology](#) website (Link 28a).

Management Option	Recommendations for Bean Rust
<b>Scouting/thresholds</b>	Record the occurrence and severity of bean rust. No thresholds are available.
<b>Resistant varieties</b>	No resistant varieties are available.
<b>Crop rotation</b>	A minimum one-year rotation is recommended.
<b>Site selection</b>	Avoid areas with poor air and soil-moisture drainage.
<b>Planting</b>	Wider row spacing reduces leaf wetness and may slow epidemic development.
<b>Postharvest</b>	Incorporate infested debris immediately after harvest to hasten decomposition of the material.
<b>Sanitation</b>	Avoid walking through the crop when the leaves are wet.

## ORGANIC SNAP BEAN PRODUCTION

At the time this guide was produced, the following materials were available in New York State for managing this pest and were allowable for organic production. Listing a pest on a pesticide label does not assure the pesticide's effectiveness. The registration status of pesticides can and does change. Pesticides must be currently registered with the New York State Department of Environmental Conservation (DEC) to be used legally in NY. However, pesticides meeting the federal requirements for minimum-risk (25(b)) pesticides do not require registration. Current NY pesticide registrations can be checked on the Pesticide Product, Ingredient, and Manufacturer System ([PIMS website](#) (Link 2)). ALWAYS CHECK WITH YOUR CERTIFIER before using a new product.

**Table 11.7 Labeled Products for Management of Bean Rust**

Class of Compound Product (Active ingredient)	Rate/Acre	PHI (days)	REI (hours)	Efficacy	Comments
<b>BIOLOGICALS</b>					
Double Nickel 55 Biofungicide ( <i>Bacillus amyloliquefaciens str. D747</i> )	0.25-3 lbs Foliar treatment.	0	4	?	
Double Nickel LC Biofungicide ( <i>Bacillus amyloliquefaciens str. D747</i> )	0.5-6 qts Foliar treatment.	0	4	?	
Regalia Biofungicide ( <i>Reynoutria sachalinensis</i> )	1 – 4 qts Foliar treatment.	0	4	?	Apply every 7-14 days as needed.
Serenade ASO ( <i>Bacillus subtilis</i> )	2-6 qts Foliar treatment.	0	4	?	
Serenade MAX ( <i>Bacillus subtilis</i> )	1-3 lbs Foliar treatment.	0	4	?	
<b>OILS</b>					
Cinerate (cinnamon oil)	13-30 fl oz/100 gal water Foliar treatment.	-	-	?	
<b>SULFUR</b>					
Micro Sulf (sulfur)	7 lbs Foliar treatment.	-	24	3	Elemental sulfur not effective in 2/2 trials. Consult processor before using sulfur.
Microthiol Disperss (sulfur)	3-10 lbs Foliar treatment.	-	24	3	Elemental sulfur not effective in 2/2 trials. Do not apply within 2 weeks of an oil application nor at temperatures over 90 degrees.
<b>OTHER</b>					
JMS Stylet-Oil Organic JMS Stylet Oil (paraffinic oil)	3-6 qts/100 gal water Foliar treatment.	0	4	?	
Oxidate 2.0 (hydrogen dioxide, peroxyacetic acid)	32 fl oz – 1 gal/100 gal water Foliar treatment.	0	Until spray has dried	?	
PERpose Plus (hydrogen peroxide/dioxide)	1 fl oz/ gal (initial/curative) Foliar treatment.  0.25 – 0.33 fl oz/ gal water (weekly preventative) Foliar treatment.	-	Until dry	?	For initial or curative use, apply higher rate for 1 to 3 consecutive days. Then follow with weekly/preventative treatment.  For weekly or preventative treatments, apply lower rate every five to seven days. At first signs of disease, use curative rate then resume weekly preventative treatment.
PureSpray Green (petroleum oil)	0.75-1.5 gal/ 50- 100 gal water. Foliar treatment.	Up to day of harvest	4	?	
Trilogy (neem extract)	0.5-1.0% in 25-100 gal water/A Foliar treatment.	Up to day of harvest	4	?	Maximum labeled use of 2 gal/acre/application.

Efficacy: 1- effective in half or more of recent university trials, 2- effective in less than half of recent university trials, 3-not effective in any known trials, ?- not reviewed or no research available PHI = pre-harvest interval, REI = restricted-entry interval. - = pre-harvest interval isn't specified on label.

## 12. Root-Lesion Nematode, *Pratylenchus penetrans*

**Time of concern:** Before planting. Long term planning is required for sustainable management

**Key characteristics:** The root lesion nematode typically does not cause characteristic symptoms on snap bean roots. However, severely infected plants may show general chlorosis and stunting as well as a reduced root system. Infection by this nematode may predispose plants to infection and damage by other soil borne pathogens. See Cornell [fact sheet](#) for more information and photos of damage (Link 40).

Management Option	Recommendations for Root-Lesion Nematode
<b>Scouting/thresholds</b>	Use a soil bioassay with soybean to assess soil root-lesion nematode infestation levels. Or, <a href="#">submit the soil sample(s) for nematode analysis</a> at a public or private nematology lab (Link 38). See Section 4: Field Selection for more information as well as the following Cornell publications for instructions:  <a href="#">Soil Sampling for Plant-Parasitic Nematode Assessment</a> (Link 39).  <a href="#">A Soil Bioassay for the Visual Assessment of Soil Infestations of Lesion Nematode</a> (Link 40).
<b>Resistant Varieties</b>	No resistant snap bean varieties are available.
<b>Crop Rotation</b>	Root-lesion nematode has over 400 hosts including many vegetable and grain crops that are planted in rotation with snap bean thus making it difficult to manage lesion nematode strictly using crop rotation once populations have reached damaging levels. Depending on the size of the infested site, marigold varieties such as ‘Polynema’ and ‘Nemagone’ are very effective at reducing nematode populations, where marigold can be established successfully.
<b>Site Selection</b>	Assay soil for nematode infestation, if needed.
<b>Biofumigant Cover Crops</b>	Cover crops with a biofumigant effect, used as green manure, may be used for managing root-lesion nematode. It is important to note that many biofumigant crops including Sudangrass, white mustard, and rapeseed are hosts to root-lesion nematode and will increase the population until they are incorporated into the soil as a green manure, at which point their decomposition products are toxic to nematodes. Research has suggested that Sudangrass hybrid ‘Trudan 8’ can be used effectively as a biofumigant to reduce root-lesion nematode populations. Cover crops such as forage pearl millet ‘CFPM 101’ and ‘Tifgrain 102’, rapeseed ‘Dwarf Essex’, and ryegrass ‘Pennant’ are poor hosts, and thus will limit the build-up or reduce root-lesion nematode populations when used as a “standard” cover crop.
<b>Sanitation</b>	Avoid moving soil from infested fields to un-infested fields via equipment and vehicles, etc. Also limit/avoid surface run-off from infested fields.
<b>Weed control</b>	Many common weed species including lambsquarters, redroot pigweed, common purslane, common ragweed, common dandelion and wild mustard are also hosts therefore effective weed management is also important.

## 13. INSECT MANAGEMENT

Effective insect management relies on accurate identification of pests and beneficial insects, an understanding of their biology and life cycle, knowledge of economically important levels of pest damage, and a familiarity with the effectiveness of allowable control practices, in other words, Integrated Pest Management (IPM).

Regular scouting and accurate pest identification are essential for effective insect management. Thresholds used for conventional production may not be useful for organic systems because of the typically lower percent mortality and shorter residual of control products allowed for organic production. The use of pheromone traps or other monitoring and prediction techniques can provide an early warning for pest problems, and help effectively focus scouting efforts.

The contribution of crop rotation as an insect management strategy is highly dependent on the mobility of the pest. Crop rotation tends to make a greater impact on reducing pest populations if the pest has limited mobility. In cases where the insects are highly mobile, leaving a greater distance between past and present plantings is better.

### Natural Enemies

Learn to identify naturally occurring beneficial insects, and attract and conserve them in your fields by providing a wide variety of flowering plants in or near the field and by avoiding use of broad-spectrum insecticides during periods when natural enemies are present. In most cases, a variety of natural enemies are present in the field, each helping to reduce pest populations. The additive effects of multiple species of natural enemies, attacking different host stages, is more likely to make an important contribution to reducing pest populations than individual natural enemy species operating alone. Natural enemies need a reason to be present in the field, either a substantial pest population, alternative hosts, or a source of pollen or nectar, and may not respond to a buildup of pests quickly enough to keep pest populations below damaging levels. Releasing insectary-reared beneficial organisms into the crop early in the pest outbreak may help control some pests but sometimes these biocontrol agents simply leave the area. For more information, see Cornell's [Natural Enemies of Vegetable Insect Pests](#) (Reference 3) and

[Biological Control: A Guide to Natural Enemies in North America](#) (Link 41).

### Regulatory

Organic farms must comply with all other regulations regarding pesticide applications. See Section 10 for details. ALWAYS check with your organic farm certifier when planning pesticide applications.

### Efficacy

In general, insecticides allowed for organic production kill a smaller percentage of the pest population and have a shorter residual than non-organic insecticides. University-based efficacy testing is not available for many organic pesticides. See Section 10.3 for more information on application techniques that can optimize effectiveness.

### Resources:

[Natural enemies of Vegetable Insect Pests](#) (Reference 3)

[Biological Control: A Guide to Natural Enemies in North America](#) (Link 41).

[Resource guide for Organic Insect and Disease Management](#) (Reference 1)

At the time this guide was produced, the following materials were available in New York State for managing this pest and were allowable for organic production. Listing a pest on a pesticide label does not assure the pesticide's effectiveness. The registration status of pesticides can and does change. Pesticides must be currently registered with the New York State Department of Environmental Conservation (DEC) to be used legally in NY. However, pesticides meeting the federal requirements for minimum-risk (25(b)) pesticides do not require registration. Current NY pesticide registrations can be checked on the Pesticide Product, Ingredient, and Manufacturer System ([PIMS website](#)) (Link 2). ALWAYS CHECK WITH YOUR CERTIFIER before using a new product.

**Table 13.0 Pesticides Labeled for Organic Bean Insect Control**

	European Corn Borer	Mexican Bean Beetle	Potato Leafhopper	Seedcorn Maggot	Two-Spotted Mite	Tarnished Plant Bug
<b>BOTANICALS</b>						
Aza-Direct ( <i>azadirachtin</i> )	X	X	X	X	X	X
AzaGuard ( <i>azadirachtin</i> )		X	X		X	X
AzaMax ( <i>azadirachtin</i> )	X	X	X		X	X
AzaSol ( <i>azadirachtin</i> )		X				
Azatrol EC ( <i>azadirachtin</i> )	X	X	X		X	X
Azera ( <i>azadirachtin and pyrethrins</i> )	X	X	X		X	X
BioLink ( <i>garlic juice</i> )	X	X	X		X	X
BioLink Insect & Bird Repellant ( <i>garlic juice</i> )	X	X	X		X	X
Ecozin Plus 1.2% ME ( <i>azadirachtin</i> )	X	X	X	X		X

**Table 13.0 Pesticides Labeled for Organic Bean Insect Control**

	European Corn Borer	Mexican Bean Beetle	Potato Leafhopper	Seedcorn Maggot	Two-Spotted Mite	Tarnished Plant Bug
Envirepel 20 ( <i>garlic juice</i> )	X	X	X		X	X
Garlic Barrier ( <i>garlic juice</i> )	X	X	X		X	X
Molt-X ( <i>azadirachtin</i> )	X	X	X			X
Neemazad 1% EC ( <i>azadirachtin</i> )		X	X			
Neemix 4.5 ( <i>azadirachtin</i> )		X	X			
PyGanic Crop Protection 1.4EC <sub>II</sub> ( <i>pyrethrin</i> )	X	X	X		X	X
PyGanic Crop Protection 5.0 EC <sub>II</sub> ( <i>pyrethrin</i> )	X	X	X		X	X
Safer Brand #567 ( <i>pyrethrin &amp; potassium salts of fatty acids</i> )		X	X		X	X
<b>BIOLOGICALS</b>						
Deliver ( <i>Bacillus thuringiensis subsp. Kurstaki</i> )	X					
DiPel DF ( <i>Bacillus thuringiensis subsp. Kurstaki</i> )	X					
Entrust Naturalyte Insect Control ( <i>spinosad</i> )	X					
Entrust SC ( <i>spinosad</i> )	X					
Grandevo ( <i>Chromobacterium subtsugae str. PR4A4-1</i> )			X		X	
Javelin ( <i>Bacillus thuringiensis subsp. Kurstaki</i> )	X					
Mycotrol O ( <i>Beauveria bassiana str. GHA</i> )	X		X			X
PFR-97 20% WDG ( <i>Isaria fumosorosea Apopka str. 97</i> )					X	X
<b>OILS</b>						
BioRepel Natural Insect Repellent ( <i>garlic oil</i> )			X			
Cedar Gard ( <i>cedar oil</i> )	X	X	X			X
Cinnerate ( <i>cinnamon oil</i> )					X	
Ecotec ( <i>rosemary and peppermint oils</i> )	X	X			X	X
GC-Mite ( <i>cottonseed, clove and garlic oils</i> )					X	
Glacial Spray Fluid ( <i>mineral oil</i> )		X	X		X	
Golden Pest Spray Oil ( <i>soybean oil</i> )		X	X		X	X
JMS Stylet Oil ( <i>paraffinic oil</i> )			X		X	
Oleotrol-I Bio-Insecticide Concentrate ( <i>soybean oil</i> )					X	
Omni Supreme Spray ( <i>mineral oil</i> )					X	
Organocide 3-in-1 Garden Spray ( <i>sesame oil</i> )					X	
PureSpray Green ( <i>petroleum oil</i> )		X	X		X	
SuffOil-X ( <i>petroleum oil</i> )		X	X		X	
TriTek ( <i>mineral oil</i> )		X	X		X	
<b>OTHER</b>						
Micro Sulf ( <i>sulfur</i> )					X	
Microthiol Disperss ( <i>sulfur</i> )					X	

**Table 13.0 Pesticides Labeled for Organic Bean Insect Control**

	European Corn Borer	Mexican Bean Beetle	Potato Leafhopper	Seedcorn Maggot	Two-Spotted Mite	Tarnished Plant Bug
M-pede ( <i>potassium salts of fatty acids</i> )			X		X	X
Nuke Em Natural Insecticide and Fungicide ( <i>citric acid</i> )					X	
Sil-Matrix ( <i>potassium silicate</i> )					X	
Surround WP Crop Protectant ( <i>kaolin</i> )		X	X			
Trilogy ( <i>neem extract</i> )					X	

X-labeled for use in NYS and also listed on the Organic Materials Review Institute

### 13.1 Mexican Bean Beetle

**Time for concern:** June through September

**Key characteristics:** Adults are 1/4-1/3 inch long, convex, and oval /in form. They vary in color from yellow when newly emerged to a coppery brown when mature. Each wing has eight black dots in three rows across the back. Eggs are orange to yellow in color and deposited in groups of 40 to 50 on the underside of leaves. Larvae are yellow, and the bodies are covered with six rows of long, black-tipped spines. Feeding by adults and larvae results in the skeletonizing of leaves. See Cornell [fact sheet](#) and [photo of damage](#) (Links 42, 43).

Management Option	Recommendations for Mexican Bean Beetle
<b>Scouting/thresholds</b>	Overwintering adults move into fields and feed for 1-2 weeks and then lay their eggs. Monitor fields by scouting for adults, eggs, and larvae. Because populations vary within a field and between fields in an area, it is difficult to determine when population numbers present a threat.
<b>Natural Enemies</b>	Natural enemies such as parasitic flies, wasps, and predators help to control Mexican bean beetle populations. An imported parasitoid, <i>Pebiobus foveolatus</i> , can be important for control. The parasitoid does not overwinter successfully, so it must be reared and released each year. Use Reference 3 or see Cornell's <a href="#">Biological Control: A Guide to Natural Enemies in North America</a> for identification of natural enemies (Link 41).
<b>Trap Crop</b>	A small strip of early-planted beans in the previous year's snap bean field could be used as a trap crop to attract overwintered beetles where they can be managed with foliar applications of products listed below or tilled under after adults have completed oviposition.
<b>Resistant Varieties</b>	No resistant varieties are available.
<b>Planting Date</b>	Avoiding early plantings can reduce damage.
<b>Cultural</b>	Avoid planting sequential crops adjacent to each other. Rotating fields as far away as possible from the previous season's fields should help to reduce populations.
<b>Postharvest</b>	Soon after harvest, plow under the infested crop to prevent immature beetles from completing development on the foliage and to destroy potential overwintering beetles.

# ORGANIC SNAP BEAN PRODUCTION

At the time this guide was produced, the following materials were available in New York State for managing this pest and were allowable for organic production. Listing a pest on a pesticide label does not assure the pesticide's effectiveness. The registration status of pesticides can and does change. Pesticides must be currently registered with the New York State Department of Environmental Conservation (DEC) to be used legally in NY. However, pesticides meeting the federal requirements for minimum-risk (25(b)) pesticides do not require registration. Current NY pesticide registrations can be checked on the Pesticide Product, Ingredient, and Manufacturer System ([PIMS website](#)) (Link 2). ALWAYS CHECK WITH YOUR CERTIFIER before using a new product.

Table 13.1 Pesticides Labeled for Management of Mexican Bean Beetle					
Class of Compound Product (Active ingredient)	Rate/A	PHI (days)	REI (hours)	Efficacy	Comments
<b>BOTANICAL</b>					
<b>Azadirachtin</b>					Azadirachtin-based products effective in 1/1 trial.
Aza-Direct	1-2 pts Foliar treatment	0	4	1	Aza-Direct may be tank mixed /at rates as low as 4 oz/A. Maximum rate is 3.5 pt/A for heavy pest infestations.
AzaGuard	8-16 fl oz Foliar treatment	0	4	1	Use in combination with OMRI approved 0.25-1% non-phytotoxic spray oil in sufficient water to cover undersides of leaves.
AzaMax	1.33 fl oz/1000 ft <sup>2</sup> Foliar treatment	0	4	1	
AzaSol	6 oz/50 gal water	-	4	1	
Azatrol EC	0.29-0.96 fl oz/1000 ft <sup>2</sup> Foliar treatment	0	4	1	
Ecozin Plus 1.2% ME	15-30 oz Foliar treatment	0	4	1	Make at least 2 applications in sequence 7-10 days apart for maximum efficacy
Molt-X	8 oz Foliar treatment	0	4	1	
Neemazad 1% EC	18-72 fl oz Foliar treatment	-	4	1	Product works on insect larvae and nymphs.
Neemix 4.5	4-16 oz Foliar treatment	-	4	1	Neemix effective in 1/1 trial. Neemix controls larvae only; apply early and often for best control.
<b>Garlic juice</b>					
BioLink	0.5-2 qts Foliar treatment	12 hr	-	?	25(b) pesticide
BioLink Insect & Bird Repellent	0.5-4 qts Foliar treatment	12 hr	-	?	25(b) pesticide
Envirepel 20	10-32 oz Foliar treatment	12 hr	12 hr	?	25(b) pesticide
Garlic Barrier AG+	See comments Foliar treatment	-	4	?	25(b) pesticide. See label for specific information.
<b>OILS</b>					
Cedar Gard (cedar oil)	1 qt Foliar treatment	-	-	?	25(b) pesticide
Ecotec (rosemary and peppermint oils)	1-4 pints/100 gal spray Foliar treatment	-	-	?	25(b) pesticide
Glacial Spray Fluid (mineral oil)	0.75-1 gal/100gal Foliar treatment	Up to day of harvest	4	?	Only for use against larvae. See label for specific application volumes
Golden Pest Spray Oil (soybean oil)	2 gal Foliar treatment	-	4	?	Only for use against larvae.

**Table 13.1 Pesticides Labeled for Management of Mexican Bean Beetle**

<b>Class of Compound Product (Active ingredient)</b>	<b>Rate/A</b>	<b>PHI (days)</b>	<b>REI (hours)</b>	<b>Efficacy</b>	<b>Comments</b>
PureSpray Green (petroleum oil)	0.75 – 1.5 gal/50- 100 gallons water Foliar treatment	Up to day of harvest	4	?	Larvae only.
SuffOil-X (petroleum oil)	1-2 gal/100 gal water Foliar treatment	Up to day of harvest	4	?	Only for use against larvae. Do not mix with sulfur products.
TriTek (mineral oil)	1-2 gal/100 gal water Foliar treatment	Up to day of harvest	4	?	Larvae only. Apply as needed.
<b>PYRETHRINS</b>					
Pyganic Crop Protection EC 1.4 II (pyrethrins)	16-64 fl oz Foliar treatment	Until dry	12	3	
Pyganic Crop Protection EC 5.0 II (pyrethrins)	4.5-17 fl oz Foliar treatment	0	12	3	
<b>OTHER</b>					
Surround WP (kaolin)	25-50 lbs Foliar treatment	Up to day of harvest	4	?	
<b>COMBINATIONS</b>					
Azera (azadirachtin, pyrethrins)	1-3 pints Foliar treatment	-	12	1	
Pyganic Crop Protection EC 1.4 II <b>OR</b> Pyganic Crop Protection EC 5.0 II (pyrethrins) <b>plus</b> Neemix 4.5 (azadirachtin)	32oz Foliar treatment  10 oz Foliar treatment  8 oz Foliar treatment	Until dry  0  -	12  12  4	1  1  1	Pyganic effective in 3/3 trials. Pyganic in combination with Neemix, provided even better protection than Pyganic alone in 1/1 trial.
Pyganic Crop Protection EC 1.4 II <b>OR</b> Pyganic Crop Protection EC 5.0 II (pyrethrins) <b>plus</b> Nu Film P (spreader sticker)	32 oz Foliar treatment  10 oz Foliar treatment  4-6 oz Foliar treatment	Until dry  0  	12  12  12	1  1  1	Pyganic in combination with Nufilm provided better protection than Pyganic alone in 1/1 trial.
Safer Brand #567 Pyrethrin & Insecticidal Soap Concentrate II (pyrethrin & potassium salts of fatty acids)	1 gal of mixed spray/700 ft <sup>2</sup> of plant surface area Foliar treatment	Until spray has dried	12	?	See label for specific mixing instructions.

Efficacy: 1- effective in half or more of recent university trials, 2- effective in less than half of recent university trials, 3-not effective in any known trials, ?- not reviewed or no research available PHI = pre-harvest interval, REI = restricted-entry interval. - = pre-harvest interval isn't specified on label.

### 13.2 Potato Leafhopper (PLH), *Empoasca fabae*

**Time for concern:** Early June through pre-bloom

**Key characteristics:** The adult is wedge-shaped, iridescent green in color, and 1/8 inch long. The body is widest at the head. See Cornell [fact sheet](#) for photo (Link 44). Eggs are laid singly on the underside of leaves. Both adults and nymphs are very active, running forwards, backwards, or sideways. The symptoms produced by feeding have been termed “hopperburn.” The first sign of hopperburn is whitening of the veins. These areas become flaccid and yellow in color, then desiccate, turn brown, and die. Leaf curling is also very common. The entire process takes four to five days. See Cornell [photo of damage](#) (Link 45) or Reference 10.

Management Option	Recommendations for Potato Leafhopper
<b>Resistant varieties</b>	Resistance of the varieties currently grown is not known.
<b>Scouting/thresholds</b>	Potato leafhoppers migrate from southern areas each year and their time of arrival varies. Check for the presence of adult potato leafhoppers by using a sweep net or by placing yellow, sticky traps near field edges. Nymphs are best sampled by visual examination of the undersides of leaves on the lower half of the plant. Bean yields are most likely to be reduced by potato leafhoppers if damage occurs before bloom. Management should occur when a threshold is met of one nymph per trifoliate leaf or when adults exceed 100/20 sweeps. On newly emerged beans, lower densities of leafhoppers than those mentioned above may be damaging. See Reference 10.
<b>Natural enemies</b>	Although a variety of natural enemies of potato leafhoppers have been reported, their impact on infestations is not well known. Use Reference 3 or see Cornell <a href="#">Biological Control: A Guide to Natural Enemies in North America</a> (Link 41).

At the time this guide was produced, the following materials were available in New York State for managing this pest and were allowable for organic production. Listing a pest on a pesticide label does not assure the pesticide’s effectiveness. The registration status of pesticides can and does change. Pesticides must be currently registered with the New York State Department of Environmental Conservation (DEC) to be used legally in NY. However, pesticides meeting the federal requirements for minimum-risk (25(b)) pesticides do not require registration. Current NY pesticide registrations can be checked on the Pesticide Product, Ingredient, and Manufacturer System [\(PIMS\) website](#) (Link 2). ALWAYS CHECK WITH YOUR CERTIFIER before using a new product.

**Table 13.2 Pesticides Labeled for Management of Potato Leafhopper**

Class of Compound Product (Active ingredient)	Rate/A	PHI (days)	REI (hours)	Efficacy	Comments
<b>BIOLOGICAL</b>					
Grandevo ( <i>Chromobacterium subtsugae</i> str. PRAA4-1)	2-3 lbs Foliar treatment	0	4	?	
Mycotrol O ( <i>Beauveria bassiana</i> strain GHA)	0.25 – 1 qt Foliar treatment	Up to day of harvest	4	?	Ground or aerial application.
<b>BOTANICAL</b>					
<b>Azadirachtin</b>					
					Azadirachtin-based product was effective in 1 /1 trial.
Aza-Direct	1-2 pints Foliar treatment	0	4	1	
AzaGuard	10-16 fl oz Foliar treatment	0	4	1	Use in combination with OMRI approved 0.25-1% non-phytotoxic spray oil in sufficient water to cover undersides of leaves.
AzaMax	1.33 fl oz/1000 ft <sup>2</sup> Foliar treatment	0	4	1	
Azatrol EC	0.24-0.96 fl oz/1000 ft <sup>2</sup> Foliar treatment	0	4	1	

**Table 13.2 Pesticides Labeled for Management of Potato Leafhopper**

<b>Class of Compound Product (Active ingredient)</b>	<b>Rate/A</b>	<b>PHI (days)</b>	<b>REI (hours)</b>	<b>Efficacy</b>	<b>Comments</b>
Ecozin Plus 1.2% ME	15-30 oz Foliar treatment	0	4	1	Make at least 2 applications in sequence 7-10 days apart for maximum efficacy
Molt-X	10 oz Foliar treatment	0	4	1	
Neemazad 1% EC	31-72 fl oz Foliar treatment	-	4	1	Product works on insect larvae and nymphs.
Neemix 4.5	7-16 oz Foliar treatment	-	4	3	Neemix alone not effective in 1/1 trial. Neemix in combination with Pyganic was effective in 1/1 trial.
<b>Garlic juice</b>					
BioLink	0.5 – 2 qts Foliar treatment	12 hr	-	?	25(b) pesticide. Repellant.
BioLink Insect & Bird Repellant	0.5-4 q/A Foliar treatment	12 hr	-	?	25(b) pesticide. Repellant.
Envirepel 20	10-32 oz/A Foliar treatment	12 hr	12 hr	?	
Garlic Barrier AG+	See comments Foliar treatment	-	4	?	25(b) pesticide. See label for specific information.
<b>Pyrethrins</b>					
Pyganic Crop Protection EC 1.4 II (pyrethrin)	16-64 fl oz Foliar treatment	Until dry	12	1	Pyganic alone effective in 2/3 trials. Two applications of Pyganic may be needed to reduce adult and nymph leafhopper populations. Pyganic in combination with Neemix effective in 1/1 trial.
Pyganic Crop Protection EC 5.0 II (pyrethrin)	4.5-17 fl oz Foliar treatment	0	12	1	
Safer Brand #567 Pyrethrin & Insecticidal Soap Concentrate II (pyrethrin & potassium salts of fatty acids)	1 gal of mixed spray/700 ft <sup>2</sup> of plant surface area Foliar treatment	-	12	?	See label for specific mixing instructions.
<b>OIL</b>					
BioRepel Natural Insect Repellent (garlic oil)	1 part product to 100 parts water Foliar treatment	-	-	?	25(b) pesticide
Cedar Gard (cedar oil)	1 qt Foliar treatment	-	-	?	25(b) pesticide
Glacial Spray Fluid (mineral oil)	0.75-1 gal/100gal Foliar treatment	Up to day of harvest	4	?	See label for specific application volumes
PureSpray Green (petroleum oil)	0.75 – 1.5 gal/50-100 gallons of water Foliar treatment	Up to day of harvest	4	?	
Golden Pest Spray Oil (soybean oil)	2 gal Foliar treatment	-	4	?	
JMS Stylet-Oil Organic JMS Stylet Oil paraffinic oil	3-6 qt/100 gal water Foliar treatment	0	4	?	
SuffOil-X (petroleum oil)	1-2 gal/100 gal water Foliar treatment	Up to day of	4	?	Do not mix with sulfur products.

**Table 13.2 Pesticides Labeled for Management of Potato Leafhopper**

Class of Compound Product (Active ingredient)	Rate/A	PHI (days)	REI (hours)	Efficacy	Comments
		harvest			
TriTek (mineral oil)	1-2 gal/100 gal water Foliar treatment	Up to day of harvest	4	?	Apply as needed.
<b>OTHER</b>					
Azera (azadirachtin, pyrethrins)	1-3 pints Foliar treatment	-	12	1	
M-pede (potassium salts of fatty acids)	0.25% - 4% volume to volume Foliar treatment	0	12	?	
Surround WP (kaolin)	25-50 lbs Foliar treatment	Up to day of harvest	4	?	

Efficacy: 1- effective in half or more of recent university trials, 2- effective in less than half of recent university trials, 3-not effective in any known trials, ?- not reviewed or no research available. PHI = pre-harvest interval, REI = restricted-entry interval. - = pre-harvest interval isn't specified on label.

### 13.3 Seedcorn Maggot, *Delia platura*

**Time for concern:** Mid-May through late June

**Key characteristics:** Adult flies resemble small house flies, are slender, 1/4 inch long, and grayish black in color. Maggots are yellowish white and infest seeds and other below-ground plant parts. See Cornell [fact sheet](#) for photo (Link 46). Maggots are only a problem from planting to plant emergence. Damaged plants are weak, may not develop, or may be delayed in maturity. Stand may be poor. See Cornell [photo of damage](#) (Link 47) or see a Cornell bulletin with [more information about damage](#) (Link 48).

Management Option	Recommendations for Seedcorn Maggot
<b>Scouting/thresholds</b>	Although there are multiple generations per year, the first generation is the important one. No thresholds are available. Five maggots per seed are required to significantly reduce stands of snap beans.
<b>Resistant varieties</b>	Snap beans are particularly susceptible.
<b>Planting date</b>	Planting after the first generation maggots have pupated will reduce damage. In New York, this occurs around June 21. Additionally, plantings after this period are less susceptible to maggot damage because warmer soil temperatures allow seedlings to emerge soon after planting, reducing the adult egg-laying period and young larval feeding period.
<b>Site selection/preparation</b>	Seedcorn maggots prefer soil with high organic matter. Incorporate crop residues at least 2 weeks before planting. Using shallow planting and other means to speed up germination and emergence will reduce damage.
<b>Seed selection/treatment</b>	No seed treatments are currently approved for organic production.
<b>Natural enemies</b>	Natural enemies can be preserved by using pesticides that are less harmful to them. Use Reference 3 see Cornell <a href="#">Biological Control: A Guide to Natural Enemies in North America</a> (Link 41).

At the time this guide was produced, the following materials were available in New York State for managing this pest and were allowable for organic production. Listing a pest on a pesticide label does not assure the pesticide's effectiveness. The registration status of pesticides can and does change. Pesticides must be currently registered with the New York State Department of Environmental Conservation (DEC) to be used legally in NY. However, pesticides meeting the federal requirements for minimum-risk (25(b)) pesticides do not require registration. Current NY pesticide registrations can be checked on the Pesticide Product, Ingredient, and Manufacturer System (PIMS) website (Link 2). ALWAYS CHECK WITH YOUR CERTIFIER before using a new product.

13.4 Labeled Products for Management of Seedcorn maggot					
Class of Compound Product (Active ingredient)	Rate/A	PHI (days)	REI (hours)	Efficacy	Comments
<b>BOTANICALS</b>					
Aza-Direct (azadirachtin)	1-2 pints Foliar spray or soil drench	0	4	?	
Ecozin Plus 1.2% ME (azadirachtin)	15-30 oz Soil drench	0	4	?	Time sprays to anticipate egg hatch or when pests first appear. Drench soil to kill larvae. Make at least 2 applications in sequence 7-10 days apart for maximum efficacy.

Efficacy: 1- effective in half or more of recent university trials, 2- effective in less than half of recent university trials, 3-not effective in any known trials, ?- not reviewed or no research available. PHI = pre-harvest interval, REI = restricted-entry interval. - = pre-harvest interval isn't specified on label

### 13.4 European Corn Borer (ECB), *Ostrinia nubilalis*

**Time for concern:** June through September.

**Key characteristics:** The eggs are laid in fish scale-like masses on the undersides of leaves. Larvae are cream colored and have a relatively smooth appearance with rows of brown spots. The head capsule is dark brown, and mature larvae are 3/4 inch long. Pupae are reddish brown, but the bean crop is harvested before pupation occurs. Adults are light yellow with reddish brown markings and are about one inch in length. See Cornell [fact sheet](#) for photo (Link 49) or Reference 2. European corn borer feeding will damage pods and their feeding within the pod will cause contamination at harvest. See Cornell [photo of damage](#) (Link 50). See reference 9.

Management Option	Recommendations for European Corn Borer
<b>Scouting/thresholds</b>	There are no formal thresholds for this pest on snap bean. Because tolerance for this pest is extremely low, insecticide use is relied upon heavily. Decision for control should be based on the stage of the bean crop and level of moth activity. Beans should be treated only when they are in the vulnerable stage, i.e., from early bud until early pod development. Moth activity can be monitored using traps (black light or pheromone) and trap catch information is updated on the Timely Tips section of the following website: <a href="#">Penn State Pest Watch</a> website (Link 51).
<b>Natural enemies</b>	A variety of natural enemies help suppress ECB populations including predatory lady beetles, minute pirate bugs and lacewings, and fly and wasp parasitoids. Natural enemies can be preserved using pesticides that are less harmful to them. Use Reference 3 or see the Cornell publication <a href="#">Biological Control: A Guide to Natural Enemies in North America</a> (Link 41). Trichogramma ostrinae, a parasitoid of eggs, has been used successfully for ECB control in sweet corn and is commercially available. Releases should be timed to susceptible crop stage and ECB flight peaks.
<b>Resistant varieties</b>	No resistant varieties are available.
<b>Crop rotation</b>	This is not a viable management option
<b>Site selection</b>	Avoid planting in or adjacent to a field that was planted with corn the previous year. However, this is likely applicable only to early-planted fields. Late-planted snap bean fields adjacent to mature untreated corn could be at risk for attack by second-generation corn borers that emerge from the mature corn.

Management Option	Recommendations for European Corn Borer
Postharvest and Sanitation	These are not currently viable management options

At the time this guide was produced, the following materials were available in New York State for managing this pest and were allowable for organic production. Listing a pest on a pesticide label does not assure the pesticide's effectiveness. The registration status of pesticides can and does change. Pesticides must be currently registered with the New York State Department of Environmental Conservation (DEC) to be used legally in NY. However, pesticides meeting the federal requirements for minimum-risk (25(b)) pesticides do not require registration. Current NY pesticide registrations can be checked on the Pesticide Product, Ingredient, and Manufacturer System ([PIMS website](#)) (Link 2). ALWAYS CHECK WITH YOUR CERTIFIER before using a new product.

### 13.4 Labeled Products for Management of European Corn Borer

Class of Compound Product (Active ingredient)	Rate/A	PHI (days)	REI (hours)	Efficacy	Comments
<b>BIOLOGICALS</b>					
Deliver ( <i>Bacillus thuringiensis subsp. kurstaki</i> )	0.25-1.5 lbs Foliar treatment	0	4	3	<i>Bacillus thuringiensis</i> products not effective in 2/2 trials. Must be eaten by the larvae to be effective; most effective against small, newly hatched larvae, so good scouting program to determine early infestations is recommended.
Javelin WG ( <i>Bacillus thuringiensis subsp. kurstaki</i> )	0.12-1.5 lbs Foliar treatment	0	4	3	
DiPel DF ( <i>Bacillus thuringiensis subsp. kurstaki</i> )	0.5-2 lbs/A Foliar treatment	0	4	3	
Entrust Naturalyte Insect Control ( <i>spinosad</i> )	1-2 oz Foliar treatment	3	4	1	Spinosad products effective in 41/47 trials. 33 studies showed good control of caterpillars, including ECB
Entrust SC ( <i>spinosad</i> )	3-6 fl oz Foliar treatment	3	4	1	Spinosad products effective in 41/47 trials. 33 studies showed good control of caterpillars, including ECB
Mycotrol O ( <i>Beauveria bassiana str. GHA</i> )	0.25 – 1 qt Foliar treatment	Up to day of harvest	4	?	Ground or aerial application
<b>BOTANICAL</b>					
Aza-Direct ( <i>azadirachtin</i> )	1-2 pints Foliar spray or soil drench	0	4	?	Can use for ECB, but requires an intensive scouting program.
AzaMax ( <i>azadirachtin</i> )	1.33 fl oz/1000 ft <sup>2</sup> Foliar treatment	0	4	?	
Azatrol EC ( <i>azadirachtin</i> )	0.24-0.96 fl oz/1000 ft <sup>2</sup> Foliar treatment	0	4	?	
Azera ( <i>azadirachtin, pyrethrins</i> )	1-3 pints Foliar treatment	-	12	?	
BioLink ( <i>garlic juice</i> )	0.5 – 2 qts Foliar treatment	12 hr	-	?	25(b) pesticide. Repellent
BioLink Insect & Bird Repellent ( <i>garlic juice</i> )	0.5 – 4 qts Foliar treatment	12 hr	-	?	25(b) pesticide. Repellent.
Ecozin Plus 1.2% ME ( <i>azadirachtin</i> )	15-30 oz Foliar spray or drench.	0	4	?	Spray soon after egg hatch. Make at least 2 applications in sequence 7-10 days apart for maximum efficacy.
Envirepel 20 ( <i>garlic juice</i> )	10-32 oz/A Foliar treatment	12 hr	12	?	25(b) pesticide. Repellent.

**13.4 Labeled Products for Management of European Corn Borer**

<b>Class of Compound Product (Active ingredient)</b>	<b>Rate/A</b>	<b>PHI (days)</b>	<b>REI (hours)</b>	<b>Efficacy</b>	<b>Comments</b>
Garlic Barrier AG+ (garlic juice)	See comments Foliar treatment	-	4	?	25(b) pesticide. See label for specific information.
Molt-X (azadirachtin)	8 oz Foliar treatment	0	4	?	
Pyganic Crop Protection EC 1.4 II (pyrethrin)	16-64 oz Foliar treatment	Until dry	12	?	
Pyganic Crop Protection EC 5.0 II (pyrethrin)	4.5-17 fl oz Foliar treatment	0	12	?	
<b>OIL</b>					
Cedar Gard (cedar oil)	1 qt Foliar treatment	-	-	?	25(b) pesticide
Ecotec (rosemary and peppermint oils)	1-4 pints/100 gal spray Foliar treatment	-	4	?	25(b) pesticide early stages only
<b>OTHER</b>					

Efficacy: 1- effective in half or more of recent university trials, 2- effective in less than half of recent university trials, 3-not effective in any known trials, ?- not reviewed or no research available. . PHI = pre-harvest interval, REI = restricted-entry interval. - = pre-harvest interval isn't specified on label

**13.5 Two-Spotted Spider Mite, *Tetranychus urticae***

**Time for concern:** June through September.

**Key characteristics:** The adult mite is yellow to dark green with two or four black, dorsal spots. See Cornell [fact sheet](#) (Link 52). Heavy [damage](#) may cause leaves to drop. Hot, dry weather favors spider mite outbreaks.

<b>Management Option</b>	<b>Recommendation for Two-Spotted Spider Mite</b>
<b>Natural enemies</b>	Natural enemies help to control spider mite populations. See the Cornell guide <a href="#">Biological Control: A Guide to Natural Enemies in North America</a> (Link 41) or use Reference 3.
<b>Resistant varieties</b>	No resistant varieties are available.
<b>Crop rotation, Site selection, and Sanitation, Postharvest</b>	These are not currently viable management options.
<b>Note(s)</b>	Dry, warm conditions may increase the chance of a two-spotted spider mite infestation.

At the time this guide was produced, the following materials were available in New York State for managing this pest and were allowable for organic production. Listing a pest on a pesticide label does not assure the pesticide's effectiveness. The registration status of pesticides can and does change. Pesticides must be currently registered with the New York State Department of Environmental Conservation (DEC) to be used legally in NY. However, pesticides meeting the federal requirements for minimum-risk (25(b)) pesticides do not require registration. Current NY pesticide registrations can be checked on the Pesticide Product, Ingredient, and Manufacturer System [\(PIMS\) website](#) (Link 2). ALWAYS CHECK WITH YOUR CERTIFIER before using a new product.

**Table 13.5 Pesticides Labeled for Management of Two-Spotted Spider Mites**

<b>Class of Compounds Product Name (active ingredient)</b>	<b>Product Rate/Acre</b>	<b>PHI (days)</b>	<b>REI (hours)</b>	<b>Efficacy</b>	<b>Comments</b>
<b>BIOLOGICALS</b>					
Grandevo ( <i>Chromobacterium subtsugae</i> str. PRAA4-1)	2-3 lbs Foliar treatment	0	4	?	
PFR-97 20% WDG ( <i>Isaria fumosorosea</i> Apopka str. 97)	1-2 lbs Foliar treatment	-	4	?	Repeat at 3-10 day intervals as needed to maintain control.

**Table 13.5 Pesticides Labeled for Management of Two-Spotted Spider Mites**

<b>Class of Compounds Product Name (active ingredient)</b>	<b>Product Rate/Acre</b>	<b>PHI (days)</b>	<b>REI (hours)</b>	<b>Efficacy</b>	<b>Comments</b>
<b>BOTANICALS</b>					
Aza-Direct ( <i>azadirachtin</i> )	1-2 pts Foliar treatment	0	4	?	
AzaGuard ( <i>azadirachtin</i> )	10-16 fl oz Foliar treatment	0	4	?	Use in combination with OMRI approved 0.25-1% non-phytotoxic spray oil in sufficient water to cover leaves.
AzaMax ( <i>azadirachtin</i> )	1.33 fl oz/1000 ft <sup>2</sup> Foliar treatment	0	4	?	
Azatrol EC ( <i>azadirachtin</i> )	0.24-0.96 fl oz/1000 ft <sup>2</sup> Foliar treatment	0	4	?	
Azera ( <i>azadirachtin, pyrethrins</i> )	1-3 pints Foliar treatment	-	12	?	
BioLink Insect Repellent ( <i>garlic juice</i> )	0.5- 2 qts Foliar treatment	12 hr	-	?	25(b) pesticide. Repellent.
BioLink Insect & Bird Repellent ( <i>garlic juice</i> )	0.5-4 q Foliar treatment	12 hr	-	?	25(b) pesticide. Repellent.
Envirepel 20 ( <i>garlic juice</i> )	10-32 oz Foliar treatment	12 hr	12 hr	?	25(b) pesticide. Repellent.
Garlic Barrier AG+ ( <i>garlic juice</i> )	See comments Foliar treatment	-	4	?	25(b) pesticide. See label for specific information.
PyGanic Crop Protection EC 1.4 II ( <i>pyrethrins</i> )	16-64 fl oz Foliar treatment	until dry	12	?	
PyGanic Crop Protection EC 5.0 II ( <i>pyrethrins</i> )	4.5 – 17 fl oz Foliar treatment	0	12	?	
Safer Brand #567 Pyrethrin & Insecticidal Soap Concentrate II ( <i>pyrethrin &amp; potassium salts of fatty acids</i> )	1 gal spray mix/700 ft <sup>2</sup> plant surface area Foliar treatment	-	12	?	See label for specific mixing instructions.
Trilogy ( <i>neem extract</i> )	0.5-2.0% in 25-100 gal water/A Foliar treatment	Up to day of harvest	4	?	Maximum labeled use of 2 gal/acre/application.
<b>OILS</b>					
Cinnerate ( <i>cinnamon oil</i> )	13-30 fl oz/ 100 gal water Apply 100-160 gal of spray/A Foliar treatment	-	-	?	
Ecotec ( <i>rosemary and peppermint oils</i> )	1-4 pints/100 gal spray Foliar treatment	-	-	?	25(b) pesticide
GC-Mite ( <i>cottonseed, clove and garlic oils</i> )	1 gal/100 gal spray water Foliar treatment	-	-	?	25(b) pesticide
Glacial Spray Fluid ( <i>mineral oil</i> )	0.75-1 gal/100gal Foliar treatment	Up to day of harvest	4	?	See label for specific application volumes
Golden Pest Spray Oil ( <i>soybean oil</i> )	2 gal Foliar treatment	-	4	?	
JMS Stylet-Oil Organic JMS Stylet Oil ( <i>paraffinic oil</i> )	3-6 qt/100 gal water Foliar treatment	0	4	?	

**Table 13.5 Pesticides Labeled for Management of Two-Spotted Spider Mites**

<b>Class of Compounds Product Name (active ingredient)</b>	<b>Product Rate/Acre</b>	<b>PHI (days)</b>	<b>REI (hours)</b>	<b>Efficacy</b>	<b>Comments</b>
Oleotrol-I Bio-Insecticide Concentrate (soybean oil)	1 part concentrate:300 parts water Foliar treatment	0	-	?	
Omni Supreme Spray (mineral oil)	1 – 2 gal/100 gallons of water Foliar treatment	-	12	?	
Organocide 3-in-1 Garden Spray (sesame oil)	2 oz/gal water Foliar treatment	-	-	?	25(b) pesticide
PureSpray Green (petroleum oil)	0.75 – 1.5 gal/50-100 gallons of water Foliar treatment	Up to day of harvest	4	?	
SuffOil-X (petroleum oil)	1-2 gal/100 gal water Foliar treatment	Up to day of harvest	4	?	Do not mix with sulfur products.
TriTek (mineral oil)	1-2 gal/100 gal water Foliar treatment	Up to day of harvest	4	?	Apply as needed.
<b>OTHER</b>					
Micro Sulf (sulfur)	7 lbs Foliar treatment	-	24	?	Consult processor before using sulfur.
Microthiol Disperss (sulfur)	3-10 lbs Foliar treatment	-	24	?	Consult processor before using sulfur.
M-pede (potassium salts of fatty acids)	0.25% - 4% volume to volume Foliar treatment	0	12	?	
Nuke Em Natural Insecticide and Funigicide (citric acid)	1 fl oz/31 oz water to 2 fl oz/30 fl oz water Foliar treatment	0	-	?	
Sil-Matrix (potassium silicate)	0.5% - 1% solution Foliar treatment	0	4	?	Mix 2-4 qts in 100 gallons of water and apply at 20 gallons finished spray/A.

Efficacy: 1- effective in half or more of recent university trials, 2- effective in less than half of recent university trials, 3-not effective in any known trials, ?- not reviewed or no research available. PHI = pre-harvest interval, REI = restricted-entry interval. - = pre-harvest interval isn't specified on label

### 13.6 Tarnished Plant Bug (TPB), *Lygus lineolaris*

**Time for concern:** June through September

**Key characteristics:** The adult is an oval, brown bug, mottled with various shades of reddish and yellowish brown, and about 1/4 inch long. The eggs are elongated and curved with a square outer end. Nymphs are small and greenish yellow. See Cornell [fact sheet](#) (Link 53). The nymphs feed more than the adults. The sucking injury from the nymphs causes buds to drop, pods to be misshapen, and plants to be stunted and distorted. Tarnished plant bug can be a problem from bloom through harvest. See Cornell [photo of damage](#) (Link 54) or Reference 8.

<b>Management Option</b>	<b>Recommendations for Tarnished Plant Bug</b>
<b>Scouting/thresholds</b>	Check for TPB on pigweed seed heads or by using a sweep net. Trials on snap beans indicated no yield reductions occurred with five adult TPBs or less per plant at blossom through pin pod stages.

<b>Natural enemies</b>	Natural enemies that can help control TPB populations can be preserved by using pesticides that are less harmful to them. See the Cornell guide <a href="#">Biological Control: A Guide to Natural Enemies in North America</a> (Link 41) or use Reference 3.
<b>Resistant varieties</b>	No resistant varieties are available. Although the TPB will feed on bean pods under New York's growing conditions, most snap bean varieties are somewhat tolerant with minimal pod-spotting resulting at harvest. It would be advisable to schedule plantings of susceptible varieties for early-season harvests since TPB numbers increase in bean fields as the season progresses.
<b>Crop rotation, Site selection, Postharvest, and Sanitation</b>	Effective weed management could eliminate potential hosts for tarnished plant bugs in fields and thereby minimize risk of injury to the snap bean crop. TPB has a wide-range of hosts including many weeds and is particularly attracted to flower buds. Minimizing weeds that tend to bloom prior to bean blooms can help reduce the overall population of TPB in the field. Avoid situations in which snap beans, in a vulnerable stage, are near hayfields where TPB numbers may build up and move into beans when the hay is cut.

At the time this guide was produced, the following materials were available in New York State for managing this pest and were allowable for organic production. Listing a pest on a pesticide label does not assure the pesticide's effectiveness. The registration status of pesticides can and does change. Pesticides must be currently registered with the New York State Department of Environmental Conservation (DEC) to be used legally in NY. However, pesticides meeting the federal requirements for minimum-risk (25(b)) pesticides do not require registration. Current NY pesticide registrations can be checked on the Pesticide Product, Ingredient, and Manufacturer System ([PIMS website](#)) (Link 2). ALWAYS CHECK WITH YOUR CERTIFIER before using a new product.

Table 13.6 Pesticides Labeled for Management of Tarnished Plant Bug					
Class of Compounds Product Name (active ingredient)	Product Rate/Acre	PHI (days)	REI (hours)	Efficacy	Comments
<b>BIOLOGICALS</b>					
Mycotrol O ( <i>Beauveria bassiana</i> str. GHA)	0.25 qt/A Foliar treatment	Up to day of harvest	4	?	Ground or aerial spray.
PFR-97 20% WDG ( <i>Isaria fumosorosea</i> Apopka str. 97)	1-2 lbs Foliar treatment	-	4	?	Repeat at 3-10 day intervals as needed to maintain control.
<b>BOTANICALS</b>					
Aza-Direct ( <i>azadirachtin</i> )	1-2 pts Foliar treatment	0	4	?	
AzaGuard ( <i>azadirachtin</i> )	10-16 fl oz Foliar treatment	0	4	?	Spray nymphs early.
AzaMax ( <i>azadirachtin</i> )	1.33 fl oz/1000 ft <sup>2</sup> Foliar treatment	0	4	?	
Azatrol EC ( <i>azadirachtin</i> )	0.24-0.96 fl oz/1000 ft <sup>2</sup> Foliar treatment	0	4	?	
Azera ( <i>azadirachtin</i> , pyrethrins)	1-3 pints Foliar treatment	-	12	?	
BioLink ( <i>garlic juice</i> )	0.5 – 2 qts Foliar treatment	12 hr	-	?	25(b) pesticide. Repellent.
BioLink Insect & Bird Repellent ( <i>garlic juice</i> )	0.5-4 q Foliar treatment	12 hr	-	?	25(b) pesticide. Repellent.
Ecozin Plus 1.2% ME ( <i>azadirachtin</i> )	15-30 oz Foliar treatment	0	4	?	Spray nymphs early and repeat application after 7 days. Make at least 2 applications in sequence maximum efficacy.

**Table 13.6 Pesticides Labeled for Management of Tarnished Plant Bug**

<b>Class of Compounds Product Name (active ingredient)</b>	<b>Product Rate/Acre</b>	<b>PHI (days)</b>	<b>REI (hours)</b>	<b>Efficacy</b>	<b>Comments</b>
Envirepel 20 (garlic juice)	10-32 oz/A Foliar treatment	12 hr	12	?	25(b) pesticide. Repellant.
Garlic Barrier AG+ (garlic juice)	See comments Foliar treatment	-	4	?	25(b) pesticide. See label for specific information.
Molt-X (azadirachtin)	10 oz Foliar treatment	0	4	?	
PyGanic Crop Protection EC 1.4 II (pyrethrins)	16 – 64 fl oz Foliar treatment	Until dry	12	?	
PyGanic Crop Protection EC 5.0 II (pyrethrins)	4.5 – 17 fl oz Foliar treatment	0	12	?	
Safer Brand #567 Pyrethrin & Insecticidal Soap Concentrate II (pyrethrin & potassium salts of fatty acids)	1 gal spray mix/700 ft <sup>2</sup> plant surface area Foliar treatment	-	12	?	See label for specific mixing instructions.
<b>OILS</b>					
Cedar Gard (cedar oil)	1 qt Foliar treatment	-	-	?	25(b) pesticide
Ecotec (rosemary and peppermint oils)	1-4 pints/100 gal spray Foliar treatment	-	-	?	25(b) pesticide
Golden Pest Spray Oil (soybean oil)	2 gal Foliar treatment	-	4	?	Only for use against larvae.
<b>OTHER</b>					
M-pede (potassium salts of fatty acids)	0.25% - 4% volume to volume Foliar treatment	0	12	?	

Efficacy: 1- effective in half or more of recent university trials, 2- effective in less than half of recent university trials, 3-not effective in any known trials, ?- not reviewed or no research available. . PHI = pre-harvest interval, REI = restricted-entry interval. - = pre-harvest interval isn't specified on label

## 14. SLUGS

**Time of concern:** May through September.

**Key characteristics:** Adult slugs are between one and two inches in length. See Cornell [fact sheet](#) (Link 55). Slugs can overwinter at any stage of development. Although slugs cannot survive prolonged subzero temperatures or desiccation, the burrows of small mammals and worms provide insulation from the weather. Slugs begin to move, hatch, feed, and lay eggs in the spring when temperatures are consistently above 40°F. There is often little or no slug activity in the field during periods of dry weather; however, there may be extensive feeding when the weather is damp.

<b>Management Option</b>	<b>Recommendations for Management of Slugs</b>
<b>Scouting/thresholds</b>	Record the occurrence and severity of slug damage. No thresholds have been established.
<b>Resistant varieties</b>	No resistant varieties are available.
<b>Cultural</b>	Practices that help dry the soil surface for example conventional tillage, good weed control, and using raised beds that dry out more readily than flat beds, will reduce slug populations. Heavy organic mulch creates an ideal environment for slugs.

At the time this guide was produced, the following materials were available in New York State for managing this pest and were allowable for organic production. Listing a pest on a pesticide label does not assure the pesticide's effectiveness. The registration status of pesticides can and does change. Pesticides must be currently registered with the New York State Department of Environmental Conservation (DEC) to be used legally in NY. However, pesticides meeting the federal requirements for minimum-risk (25(b)) pesticides do not require registration. Current NY pesticide registrations can be checked on the Pesticide Product, Ingredient, and Manufacturer System ([PIMS website](#) (Link 2)). ALWAYS CHECK WITH YOUR CERTIFIER before using a new product.

**Table 14. Pesticides Labeled for Management of Slugs**

Class of Compound Product (Active ingredient)	Rate/A	PHI (days)	REI (hours)	Efficacy	Comments
<b>IRON</b>					
Sluggo AG (iron phosphate)	20-44 lbs Soil treatment	0	0	?	Treat field perimeter. Lasts up to 4 weeks.
Sluggo Slug and Snail Bait (iron phosphate)	20-44 lbs Soil treatment	-	0	?	Scatter bait around the perimeter of the vegetable- plantings.
<b>IRON AND SPINOSAD</b>					
Bug-N-Sluggo (iron phosphate and spinosad)	20-44 lbs Soil treatment	3	4	?	
<b>BOTANICAL</b>					
BioLink (garlic juice)	0.5 - 2 qts Foliar treatment	12 hr	-	?	25(b) pesticide. Repellant.
BioLink Insect & Bird Repellant (garlic juice)	0.5-4 q/A Foliar treatment	12 hr	-	?	25(b) pesticide. Repellant.

Efficacy: 1- effective in half or more of recent university trials, 2- effective in less than half of recent university trials, 3-not effective in any known trials, ?- not reviewed or no research available. PHI = pre-harvest interval, REI = restricted-entry interval. - = pre-harvest interval isn't specified on label

## 15. PESTICIDES & ABBREVIATIONS MENTIONED IN THIS PUBLICATION

**Table 1. Insecticides and molluscicides mentioned in this publication**

TRADE NAME	COMMON NAME	EPA REG. NO.
Aza-Direct	<i>azadirachtin</i>	71908-1-10163
AzaGuard	<i>azadirachtin</i>	70299-17
AzaMax	<i>azadirachtin</i>	71908-1-81268
AzaSol	<i>azadirachtin</i>	81899-4
Azatrol EC	<i>azadirachtin</i>	2217-836
Azera	<i>azadirachtin, pyrethrins</i>	1021-1872
BioLink	<i>garlic juice</i>	exempt-25(b) pesticide
BioLink Insect & Bird Repellant	<i>garlic juice</i>	exempt 25(b) pesticide
BioRepel Natural Insect Repellent	<i>garlic oil</i>	exempt-25(b) pesticide
Bug-N-Sluggo	<i>iron phosphate and spinosad</i>	67702-24-70051
Cedar Gard	<i>cedar oil</i>	exempt-25(b) pesticide
Cinerate	<i>cinnamon oil</i>	exempt-25(b) pesticide
Deliver	<i>Bacillus thuringiensis subsp. kurstaki</i>	70051-69
DiPel DF	<i>Bacillus thuringiensis subsp. kurstaki</i>	73049-39
Ecotec	<i>rosemary and peppermint oil</i>	exempt-25(b) pesticide
Ecozin Plus 1.2% ME	<i>azadirachtin</i>	5481-559
Entrust Naturalyte Insect Control	<i>spinosad</i>	62719-282
Entrust SC	<i>spinosad</i>	62719-621

**Table 1. Insecticides and molluscicides mentioned in this publication**

TRADE NAME	COMMON NAME	EPA REG. NO.
Envirepel	<i>garlic juice</i>	exempt-25(b) pesticide
Garlic Barrier	<i>garlic juice</i>	exempt-25(b) pesticide
GC-Mite	<i>cottonseed, clove and garlic oils</i>	exempt-25(b) pesticide
Glacial Spray Fluid	<i>mineral oil</i>	34704-849
Golden Pest Spray Oil	<i>soybean oil</i>	57538-11
Grandevo	<i>Chromobacterium subtsugae str. PRAA4-1</i>	84059-17
Javelin WG	<i>Bacillus thuringiensis subsp. kurstaki</i>	70051-66
JMS Stylet Oil	<i>paraffinic oil</i>	65564-1
Micro Sulf	<i>sulfur</i>	55146-75
Molt-X	<i>azadirachtin</i>	68539-11
M-pede	<i>potassium salts of fatty acids</i>	10163-324
Mycotrol O	<i>Beauveria bassiana</i>	82074-3
Neemazad 1%EC	<i>azadirachtin</i>	70051-104
Neemix 4.5	<i>azadirachtin (neem)</i>	70051-9
Nuke Em Natural Insecticide and Fungicide	<i>citric acid</i> <i>soybean oil</i>	exempt-25(b) pesticide
Oleotrol-I Bio-Insecticide Concentrate	<i>mineral oil</i>	exempt-25(b) pesticide
Omni Supreme Spray		5905-368
Organic JMS Stylet Oil	<i>paraffinic oil</i>	65564-1
Organocide 3-in-1 Garden Spray	<i>sesame oil</i>	exempt-25(b) pesticide
PureSpray Green	<i>petroleum oil</i>	69526-9
PFR-97 20% WDG	<i>Isaria fumosorosea Apopka str. 97</i>	70051-19
Pyganic Crop Protection EC 1.4 <sub>II</sub>	<i>pyrethrin</i>	1021-1771
Pyganic Crop Protection EC 5.0 <sub>II</sub>	<i>pyrethrin</i>	1021-1772
Safer Brand #567 Pyrethrin & Insecticidal Soap Concentrate II	<i>pyrethrin &amp; potassium salts of fatty acids</i>	59913-9
Sil-Matrix	<i>potassium silicate</i>	82100-1
SuffOil-X	<i>petroleum oil</i>	48813-1-68539
Sluggo AG	<i>iron phosphate</i>	67702-3-54705
Sluggo Slug & Snail Bait	<i>iron phosphate</i>	67702-3-70051
Surround WP	<i>kaolin</i>	61842-18
Trilogy	<i>neem extract</i>	70051-2
TriTek	<i>mineral oil</i>	48813-1

**Table 2. Fungicides mentioned in this publication**

TRADE NAME	COMMON NAME	EPA REG. NO.
Actinovate AG	<i>Streptomyces lydicus WYEC 108</i>	73314-1
Actinovate STP	<i>Streptomyces lydicus WYEC 108</i>	73314-4
Badge X2	<i>Copper oxychloride, Copper hydroxide</i>	80289-12
Basic Copper 53	<i>basic copper sulfate</i>	45002-8
BIO-TAM	<i>Trichoderma asperellum,</i> <i>Trichoderma gamsii</i>	80289-9-69592
Champ WG	<i>copper hydroxide</i>	55146-1
Cueva Fungicide Concentrate	<i>copper octanoate</i>	67702-2-70051
Contans WG	<i>Coniothyrium minitans</i>	72444-1
CS 2005	<i>copper sulfate pentahydrate</i>	66675-3
Double Nickel 55 Biofungicide	<i>Bacillus amyloliquefaciens str. D747</i>	70051-108
Double Nickel LC Biofungicide	<i>Bacillus amyloliquefaciens str. D747</i>	70051-107
EcoMate ARMICARB O	<i>potassium bicarbonate</i>	5905-541
JMS Stylet Oil	<i>paraffinic oil</i>	65564-1

**Table 2. Fungicides mentioned in this publication**

TRADE NAME	COMMON NAME	EPA REG. NO.
Micro Sulf	<i>sulfur</i>	55146-75
Microthiol Disperss	<i>sulfur</i>	70506-187
Mycostop Biofungicide	<i>Streptomyces griseoviridis</i>	64137-5
Mycostop Mix	<i>Streptomyces griseoviridis</i>	64137-9
Nordox 75 WG	<i>cuprous oxide</i>	48142-4
NuCop 50DF	<i>copper hydroxide</i>	45002-4
Nu-Cop HB	<i>copper hydroxide</i>	42750-132
Nu-Cop 50 WP	<i>copper hydroxide</i>	45002-7
Optiva	<i>Bacillus subtilis</i> str. QST 713	69592-26
Organic JMS Stylet Oil	<i>paraffinic oil</i>	65564-1
Oxdate 2.0	<i>hydrogen dioxide</i>	70299-12
PERpose Plus	<i>hydrogen peroxide/dioxide</i>	86729-1
Prestop Biofungicide Powder	<i>Gliocladium catenulatum</i> str. J1446	64137-11
PureSpray Green	<i>petroleum oil</i>	69526-9
Regalia Biofungicide	<i>Reynoutria sachalinensis</i>	84059-3
RootShield Granules	<i>Trichoderma</i>	68539-3
RootShield PLUS+ Granules	<i>Trichoderma species</i>	68539-10
RootShield PLUS+ WP	<i>Trichoderma species</i>	68539-9
Serenade ASO	<i>Bacillus subtilis</i>	69592-12 and 264-1152
Serenade MAX	<i>Bacillus subtilis</i>	69592-11 and 264-1151
Serenade Optimum	<i>Bacillus subtilis</i>	264-1160
Serenade Soil	<i>Bacillus subtilis</i>	69592-12 and 264-1152
SoilGard Microbial Fungicide	<i>Gliocladium virens</i> str. GL-21	70051-3
Taegro	<i>Bacillus subtilis</i>	70127-5
TerraClean 5.0	<i>hydrogen dioxide, peroxyacetic acid</i>	70299-13
Trilogy	<i>neem oil</i>	70051-2
Zonix	<i>Rhamnolipid Biosurfactant</i>	72431-1

**Abbreviations and Symbols Used in This Publication**

A acre	NE not effective
AG agricultural use label	NI no information
AR annual rye	NFT not frost tolerant
ASO aqueous suspension-organic	P phosphorus
AS aqueous suspension	PHI pre-harvest interval
DF dry flowable	P <sub>2</sub> O <sub>5</sub> phosphorus oxide
EC emulsifiable concentrate	PR perennial rye
F flowable	R resistant varieties
HC high concentrate	REI restricted entry interval
K potassium	WP wettable powder
K <sub>2</sub> O potassium oxide	WG water dispersible granular
N nitrogen	WPS Worker Protection Standard

**16. REFERENCES**

- 1 Caldwell, B. Rosen, E. B., Sideman, E., Shelton, A. M., Smart, C. (2013). *Resource Guide for Organic Insect and Disease Management 2<sup>nd</sup> Ed.* New York State Agricultural Experiment Station, Geneva, NY. (<http://web.pppmb.cals.cornell.edu/resourceguide/pdf/resource-guide-for-organic-insect-and-disease-management.pdf>).
- 2 Colorado State University. (1996). *Dry Bean Production and Pest Management*. Regional Bulletin 562A, Cooperative Extension Resource Center, 115 General Services Building, Fort Collins, CO.

- 3 Hoffmann, M. P., and Frodsham A. C. (1993). *Natural Enemies of Vegetable Insect Pests*. (64 pp). Cornell Cooperative Extension. New York State Agricultural Experiment Station, Geneva, NY. (<http://nysaes-bookstore.myshopify.com/products/natural-enemies-of-vegetable-insect-pests>)
- 4 New York State Integrated Pest Management Program. (2008). *Integrated Crop and Pest Management Guidelines for Commercial Vegetable Production*: Chapter 13, Beans-Dry and Snap. Cornell Cooperative Extension, Geneva, NY. (<https://demo.cuguidelines.net/Guidelines/VegCrops/index26.aspx>).
- 5 Petzoldt, C. H., Pederson L. H., and Koplinka-Loehr, C. eds. (1990). *Snap Bean Pest Management: A Guide to Regular Field Monitoring in New York*. IPM Publication. 105b. New York State Agricultural Experiment Station, Geneva, NY.
- 6 Sarrantonio, M. (1994) *Northeast Cover Crop Handbook*. Rodale Institute, PA. (<http://www.amazon.com/Northeast-Cover-Crop-Handbook-Health/dp/0913107174>).
- 7 Zitter, T. A., McGrath, M. T. *Vegetable MD Online*. ([http://vegetablemdonline.ppath.cornell.edu/factsheets/Bean\\_List.htm](http://vegetablemdonline.ppath.cornell.edu/factsheets/Bean_List.htm)).
- 8 Muka, A. A. (1983). Tarnished plant bug, In *Vegetable Crops: Insects of Bean, Potato, and Celery* (p. 771.00). New York State Agricultural Experiment Station, Geneva, NY.
- 9 Showers, W. B., Witkowski J. F., Mason C. E., Calvin D. D., Higgins R. A., and Dively G. P. (1989). *European Corn Borer: Development and Management*. North Central Regional Extension Publication 327. Iowa State University, Ames, IA.
- 10 Tingey, W. M., Muka A. A. (1983). Potato leafhopper, In *Vegetable Crops: Insects of Vegetables* (p. 760.20). New York State Agricultural Experiment Station, Geneva, NY.
- 11 Stivers, L.J., Brainard, D.C. Abawi, G.S., Wolfe, D.W. (1999) *Cover Crops for Vegetable Production in the Northeast*. Cornell Cooperative Extension, Ithaca, NY (<http://ecommons.library.cornell.edu/bitstream/1813/3303/2/Cover%20Crops.pdf>).
- 12 Dillard, H. R., and Cobb, A. C. 2008. *Alternaria alternata* and *Plectosporium tabacinum* on snap beans: Pathogenicity, cultivar reaction, and fungicide efficacy. Online. Plant Health Progress doi:10.1094/PHP-2008-1212-01-RS.

## 17. WORLD WIDE WEB LINKS

All links accessed 15 April 2015

### General

- 1 United States Department of Agriculture. Agricultural Research Service. 2012 Revised Plant Hardiness Zone Map for New York ( <http://planthardiness.ars.usda.gov/PHZMWeb/>).
- 2 Pesticide Product Ingredient, and Manufacturer System (PIMS). (<http://pims.psur.cornell.edu>).

### Certification

- 3 Organic Materials Review Institute. (<https://www.omri.org/>).
- 4 New York Department of Agriculture and Markets, Organizations Providing Organic Certification Services for Producers and Processors in New York State. (<http://www.agriculture.ny.gov/AP/organic/docs/Organizations-Providing-Organic-Certification-Services.pdf>).
- 5 New York Department of Agriculture and Markets, Organic Farming Development/Assistance. (<http://www.agriculture.ny.gov/AP/organic/index.html>).
- 6 Agriculture Marketing Service, National Organic Program. (<http://www.ams.usda.gov/nop/NOP/standards/ProdHandPre.html>).
- 7 National Sustainable Agriculture Information Service, Organic Farming. (<http://attra.ncat.org/organic.html>).
- 8 Rodale Institute. (<http://www.rodaleinstitute.org/>).

### Soil Health, Cover Crops, and Crop Rotation

- 9 Björkman, Thomas. Cornell University, *Cover Crops for Vegetable Growers*. (<http://covercrops.cals.cornell.edu/decision-tool.php>).
- 10 Magdoff, F., Van Es, H., (2010). Sustainable Agriculture Research and Education, *Building Soils for Better Crops, 3rd Edition*. (<http://www.sare.org/Learning-Center/Books/Building-Soils-for-Better-Crops-3rd-Edition>),
- 11 Cornell University, Department of Horticulture. *Soil Health Website*. (<http://soilhealth.cals.cornell.edu/>).
- 11 a Mohler, C. L. and Johnson, S. E., editors. (2009). *Crop Rotation on Organic Farms: A Planning Manual*. Sustainable Agriculture Research and Education. Natural Resource, Agriculture and Engineering Service. Cooperative Extension, Ithaca NY. ([http://www.nraes.org/nra\\_crof.html](http://www.nraes.org/nra_crof.html)).

## Weed Management

- 12 Bowman, G., (1997). The Sustainable Agriculture Network. *Steel in the Field*. Beltsville, MD. ([http://nydairyadmin.cce.cornell.edu/uploads/doc\\_20.pdf](http://nydairyadmin.cce.cornell.edu/uploads/doc_20.pdf)).
- 13 Cornell University, Weed Ecology and Management Laboratory. (<http://weedecology.css.cornell.edu/>).
- 14 Rutgers University, *New Jersey Weed Gallery* (<http://njaes.rutgers.edu/weeds/>).
- 15 University of Vermont, Videos for Vegetable and Berry Growers. (<http://www.uvm.edu/vtvegandberry/Videos/videos.html>).
- 16 Sullivan, P., National Sustainable Agriculture Information Service (formerly ATTRA), *Principles of Sustainable Weed Management for Croplands*. (<http://attra.ncat.org/attra-pub/weed.html>).
- 17 Colquhoun, J., Bellinder, R., Cornell University. *New Cultivation Tools for Mechanical Weed Control in Vegetables*. (<http://www.vegetables.cornell.edu/weeds/newcultivationmech.pdf>).

## Crop and Soil Nutrition

- 18 Cornell Nutrient Analysis Laboratory. (<http://cnal.cals.cornell.edu/>).
- 19 Agri Analysis, Inc.. (<http://www.agrianalysis.com/>).
- 20 A&L Eastern Agricultural Laboratories, Inc. (<http://al-labs-eastern.com/>).
- 21 The Pennsylvania State University, Agricultural Analytical Services Laboratory. (<http://aasl.psu.edu>).
- 22 Dairy One Forage Lab, Ithaca, NY. (<http://dairyone.com/analytical-services/agronomy-services/soil-testing/>).
- 23 University of Massachusetts, Soil and Plant Tissue Testing Laboratory. (<http://soiltest.umass.edu/>).
- 24 Analytical Laboratory and Maine Soil Testing Service, University of Maine. (<http://anlab.umesci.maine.edu/>).
- 25 Rosen, C., Bierman, P. Using Manure and Compost as Nutrient Sources for Fruit and Vegetable Crops. University of Minnesota. (<http://www.extension.umn.edu/distribution/horticulture/M1192.html>).
- 25 a. The Pennsylvania State University. (2013). *Penn State Agronomy Guide Section 2: Soil Fertility Management*. Department of Agronomy. University Park, PA. (<http://extension.psu.edu/agronomy-guide/cm/sec2>).
- 25b Sánchez, E. S. and Richard, T. L., (2009) Pennsylvania State University Publication, UJ256. *Using Organic Nutrient Sources*. (<http://extension.psu.edu/publications/uj256>).
- 25c DuPont, T. (2011) Pennsylvania State University Publication, *Determining Nutrient Applications for Organic Vegetables*. (<http://extension.psu.edu/business/start-farming/soils-and-soil-management/determining-nutrient-applications-for-organic-vegetables-basic-calculations-introduction-to-soils-fact-3>).

## Managing Bean Diseases

- 26 Abawi, G. S., and Hunter J. E. (1979). **White Mold of Beans in New York**. (4 pp.). New York's Food and Life Sciences Bulletin 77. New York State Agricultural Experiment Station, Geneva, NY. (<http://ecommons.library.cornell.edu/bitstream/1813/5083/1/FLS-077.pdf>).
- 27 Vegetable MD online. *Photo Gallery of Important New York Vegetable diseases: Bean: White Mold*. ([http://vegetablemdonline.ppath.cornell.edu/PhotoPages/Impt\\_Diseases/Beans/Bean\\_White.htm](http://vegetablemdonline.ppath.cornell.edu/PhotoPages/Impt_Diseases/Beans/Bean_White.htm)).
- 28 George S. Abawi and J.E. Hunter (1979). Vegetable MD Online. **White Mold of Beans**. New York's Food and Life Sciences Bulletin No.77 ([http://vegetablemdonline.ppath.cornell.edu/factsheets/Beans\\_WhiteMold.htm](http://vegetablemdonline.ppath.cornell.edu/factsheets/Beans_WhiteMold.htm)).
- 28a. Dillard Lab Vegetable Pathology Website. Cornell University. College of Agriculture and Life Sciences. New York State Agricultural Experiment Station. (<http://web.pppmb.cals.cornell.edu/dillard/>).
- 29 Vegetable MD online. Photo Gallery of Important New York Vegetable diseases: **Bean: Gray mold**. ([http://vegetablemdonline.ppath.cornell.edu/PhotoPages/Impt\\_Diseases/Beans/Bean\\_Gray.htm](http://vegetablemdonline.ppath.cornell.edu/PhotoPages/Impt_Diseases/Beans/Bean_Gray.htm)).
- 30 Abawi, G. S., Crosier D. C., and Cobb A. C. (1985). **Root Rot of Snap Beans in New York**. (8 pp). New York's Food and Life Sciences Bulletin 110. New York State Agricultural Experiment Station, Geneva, NY. (<http://ecommons.library.cornell.edu/bitstream/1813/5141/1/FLS-110.pdf>).
- 31 Cornell University, Vegetable MD Online. *Photo Gallery of Important New York Vegetable Diseases: Bean: Pythium Root Rot*. ([http://vegetablemdonline.ppath.cornell.edu/PhotoPages/Impt\\_Diseases/Beans/Bean\\_Phythium.htm](http://vegetablemdonline.ppath.cornell.edu/PhotoPages/Impt_Diseases/Beans/Bean_Phythium.htm)).
- 32 Cornell University, Vegetable MD Online. *Photo Gallery of Important New York Vegetable diseases: Bean: Rhizoctonia Root Rot*. ([http://vegetablemdonline.ppath.cornell.edu/PhotoPages/Impt\\_Diseases/Beans/Bean\\_Rhizo.htm](http://vegetablemdonline.ppath.cornell.edu/PhotoPages/Impt_Diseases/Beans/Bean_Rhizo.htm)).

- 33 Dillard, H. R., and Legard, D. E. (1991). Cornell University, Vegetable MD Online. **Bacterial Diseases of Beans**. (p. 729.50). New York State Agricultural Experiment Station, Geneva, NY. ([http://vegetablemdonline.ppath.cornell.edu/factsheets/Beans\\_Bacterial.htm](http://vegetablemdonline.ppath.cornell.edu/factsheets/Beans_Bacterial.htm)).
- 34 Zitter, T.A., Provvidenti, R. (1984). Cornell University, Vegetable MD Online. **Virus Diseases of Snap and Dry Beans**. ([http://vegetablemdonline.ppath.cornell.edu/factsheets/Virus\\_Beans.htm](http://vegetablemdonline.ppath.cornell.edu/factsheets/Virus_Beans.htm)).
- 35 Cornell University, Vegetable MD Online. **Bean Virus Photo Collection**. (<http://vegetablemdonline.ppath.cornell.edu/PhotoPages/Bean/Viruses/BVphotoList.htm>).
- 36 Zitter, T. A. (2001). Cornell University Vegetable MD Online. *A Checklist of Major Weeds and Crops as Natural Hosts for Plant Viruses in the Northeast*. (<http://vegetablemdonline.ppath.cornell.edu/Tables/WeedHostTable.html>).
- 37 Mercure, P. S. (1998). University of Connecticut. **Bean Rust**. (<http://ipm.uconn.edu/documents/raw2/Bean%20Rust/Bean%20Rust.php?aid=114> ).

## Managing Nematodes

- 38 Plant Disease Diagnostic Clinic. Cornell University. (<http://plantclinic.cornell.edu>).
- 39 Abawi, F.S., Gugino, B.K. (2007) Cornell University, New York State Agricultural Experiment Station. *Soil Sampling for Plant-Parasitic Nematode Assessment*. (<http://www.fruit.cornell.edu/berrytool/pdfs/Soil%20Sampling%20for%20Nematode%20Assessment%20Factsheet.pdf> ).
- 40 Gugino, B.K., Ludwig, J.W., Abawi, G.S., Cornell University, New York State Agricultural Experiment Station. *A Soil Bioassay for the Visual Assessment of Soil Infestations of Lesion Nematode*. ([http://www.nysipm.cornell.edu/factsheets/vegetables/Lesion\\_Nematode\\_Bioassay.pdf](http://www.nysipm.cornell.edu/factsheets/vegetables/Lesion_Nematode_Bioassay.pdf) ).

## Managing Bean Insects

- 41 Weeden, C.R., Shelton, A.M., Hoffmann, M. P., *Biological Control: A Guide to Natural Enemies in North America*. (<http://www.biocontrol.entomology.cornell.edu/index.php> ).
- 42 Wilsey, W.T., Weeden, C.R., Shelton, A.M. (updated 2007). Cornell University. Pests in the Northeastern United States. **Mexican Bean Beetle**. (<http://web.entomology.cornell.edu/shelton/veg-insects-ne/pests/mbb.html> ).
- 43 Wilsey, W.T., Weeden, C.R., Shelton, A.M. (updated 2001). Cornell University. Pests in the Northeastern United States. **Mexican Bean Beetle- Damage to Bean**. ([http://web.entomology.cornell.edu/shelton/veg-insects-ne/damage/mbb\\_beans.html](http://web.entomology.cornell.edu/shelton/veg-insects-ne/damage/mbb_beans.html) ).
- 44 Wilsey, W.T., Weeden, C.R., Shelton, A.M. (updated 2001). Cornell University. Pests in the Northeastern United States. **Potato Leafhopper Life Cycle**. (<http://web.entomology.cornell.edu/shelton/veg-insects-ne/pests/plh.html> ).
- 45 Wilsey, W.T., Weeden, C.R., Shelton, A.M. (updated 2001). Cornell University. Pests in the Northeastern United States. **Potato Leafhopper Damage to Beans**. ([http://web.entomology.cornell.edu/shelton/veg-insects-ne/damage/plh\\_beans.html](http://web.entomology.cornell.edu/shelton/veg-insects-ne/damage/plh_beans.html) ).
- 46 Wilsey, W.T., Weeden, C.R., Shelton, A.M. (updated 2007). Cornell University. Pests in the Northeastern United States. **Seedcorn Maggot Life Cycle**. (<http://web.entomology.cornell.edu/shelton/veg-insects-ne/pests/scm.html> ).
- 47 Wilsey, W.T., Weeden, C.R., Shelton, A.M. (updated 2007). Cornell University. Pests in the Northeastern United States. **Seedcorn Maggot Damage to Beans**. ([http://web.entomology.cornell.edu/shelton/veg-insects-ne/damage/scm\\_beans.html](http://web.entomology.cornell.edu/shelton/veg-insects-ne/damage/scm_beans.html) ).
- 48 Vea, E. V., Webb D. R., and Eckenrode C. J. (1975). **Seedcorn Maggot Injury** (4 pp.). New York's Food and Life Sciences Bulletin 55. New York State Agricultural Experiment Station, Geneva, NY. (<http://ecommons.cornell.edu/bitstream/1813/5059/1/FLS-055.pdf> ).
- 49 Wilsey, W.T., Weeden, C.R., Shelton, A.M. (updated 2007). Cornell University. Pests in the Northeastern United States. **European Corn Borer Life Cycle**. (<http://web.entomology.cornell.edu/shelton/veg-insects-ne/pests/ecb.html> ).
- 50 Wilsey, W.T., Weeden, C.R., Shelton, A.M. (updated 2007). Cornell University. Pests in the Northeastern United States. **European Corn Borer Damage to Bean**. ([http://web.entomology.cornell.edu/shelton/veg-insects-ne/damage/ecb\\_beans.html](http://web.entomology.cornell.edu/shelton/veg-insects-ne/damage/ecb_beans.html) ).
- 51 Fleischer, S. J., Penn State, **PestWatch**. University Park, PA (<http://www.pestwatch.psu.edu/>).
- 52 Wilsey, W.T., Weeden, C.R., Shelton, A.M. (updated 2007). Cornell University. Pests in the Northeastern United States. **Twospotted Spider Mite Life Cycle**. (<http://web.entomology.cornell.edu/shelton/veg-insects-ne/pests/tsm.html> ).
- 53 Cornell Cooperative Extension. New York State Integrated Pest Management Program. (1991). **Tarnished Plant Bug**. Insect Identification Sheet No. 21, 1991 (<http://www.nysipm.cornell.edu/factsheets/treefruit/pests/tpb/tpb.asp>).
- 54 Wilsey, W.T., Weeden, C.R., Shelton, A.M. (updated 2007). Cornell University. Pests in the Northeastern United States. **Tarnished Plant Bug Damage to Beans**. ([http://web.entomology.cornell.edu/shelton/veg-insects-ne/damage/tpb\\_beans.html](http://web.entomology.cornell.edu/shelton/veg-insects-ne/damage/tpb_beans.html) ).
- 55 Wilsey, W.T., Weeden, C.R., Shelton, A.M. (updated 2007). Cornell University. Pests in the Northeastern United States. **Slugs Life Cycle**. (<http://web.entomology.cornell.edu/shelton/veg-insects-ne/pests/slugs.html> ).

## Pesticide Use, Safety, and Sprayer Calibration

- 56 Cornell Integrated Crop and Pest Management Guidelines (2009). *Chapter 6 Pesticide Information and Safety*.: (<http://veg-guidelines.cce.cornell.edu/>).
- 57 *Calibration: Backpack Sprayer*. Pesticide Environmental Stewardship. Center for Integrated Pest Management. (<http://pesticidestewardship.org/calibration/Pages/BackpackSprayer.aspx>).
- 58 Dill, J. and Koehler, G. (2005). Agricultural Pocket Pesticide Calibration Guide. University of Maine Cooperative Extension and Northeastern IPM Center. (<http://www.udel.edu/pesticide/CalibrationGuide-small.pdf>).
- 59 Landers, A., Knapsack Sprayers: General Guidelines for Use. Cornell University, Ithaca, N.Y. (<http://www.docstoc.com/docs/16502240/Scope>).
- 60 Miller, A. and Bellinder, R. (2001) Herbicide Application Using a Knapsack Sprayer. Department of Horticultural Science, Cornell University, Ithaca, N.Y. (<http://www.hort.cornell.edu/bellinder/spray/southasia/pdfs/knapsack.pdf>).
- 61 Cornell University Law School. Legal Information Institute. 40 CFR 152.25 – Exemptions for pesticides of a character not requiring FIFRA regulation. (<http://www.law.cornell.edu/cfr/text/40/152.25>).
- 62 Office of prevention, pesticides and toxic substances. (2009). *Inert ingredients eligible for FIFRA 25(b) pesticide products*. United States Environmental Protection Agency. Washington DC. ([http://www.epa.gov/opprd001/inerts/section25b\\_inerts.pdf](http://www.epa.gov/opprd001/inerts/section25b_inerts.pdf)).
- 63 Federal Insecticide Fungicide Rodenticide Act (FIFRA). (2009). *Electronic Code of Federal Regulations*. Title 7: Agriculture. National Organic Program, Part 205, sections 600-606. (<http://ecfr.gpoaccess.gov/cgi/t/text/text-idx?c=ecfr&sid=fbc697b0474ea6a90b9b31be05ddc029&rgn=div6&view=text&node=7:3.1.1.9.32.7&idno=7>).
- 64 Extension: America's Research Based Learning Network. (2015) Pesticide Environmental Stewardship Community Page. (<http://www.extension.org/pesticidestewardship>).
- 65 Center for Integrated Pest Management. Pesticide Safety Education Program, Cooperative Extension Service. Pesticide Environmental Stewardship Website. (<http://pesticidestewardship.org/Pages/About.aspx>).
- 66 Landers, Andrew. Cornell University Department of Entomology. (2003) Vegetable Spraying. (<http://web.entomology.cornell.edu/landers/pestapp/vegetable.htm>).
- 67 EPA Office of Prevention, Pesticides, and Toxic Substances. (2005). How To Comply with the Worker Protection Standard for Agricultural Pesticides: What Employers Need to Know. Unit 2: An Introduction to the Worker Protection Standard. 16 pp. ([http://www.epa.gov/agriculture/epa-735-b-05-002\\_unit2.pdf](http://www.epa.gov/agriculture/epa-735-b-05-002_unit2.pdf)).
- 68 National Pesticide Information Center: State Pesticide Regulatory Agencies. Cooperative agreement between **Oregon State University** and the U.S. Environmental Protection Agency. (<http://npic.orst.edu/mlrDetail.html?lang=en&to=SPE&state=NY#statePesticide>).
- 69 Pesticide Management Education Program (PMEP). (2013). Cornell University Cooperative Extension. (<http://psep.cce.cornell.edu/Default.aspx>).

This guide is published by the New York State Integrated Pest Management Program, which is funded through Cornell University, Cornell Cooperative Extension, the New York State Department of Agriculture and Markets, the New York State Department of Environmental Conservation, and USDA-NIFA. Copyright © 2015 New York State Integrated Pest Management Program and Cornell University. Cornell Cooperative Extension provides equal program and employment opportunities. NYS IPM Publication number 132. June 2015. [www.nysipm.cornell.edu](http://www.nysipm.cornell.edu).